

Changes in sugar content of the brain during narcosis

Gepubliceerd: 13-12-2018 Laatste bijgewerkt: 15-05-2024

As plasma glucose during surgery increases, because of the surgical stress response and accompanied insulin resistance, this will lead to an acute supply-demand mismatch and excess of glucose in the brain. We hypothesize that this leads to...

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Observationeel onderzoek, zonder invasieve metingen

Samenvatting

ID

NL-OMON20502

Bron

Nationaal Trial Register

Verkorte titel

Cerebrospinal fluid study

Aandoening

cerebral glucose metabolism; polyol pathway; Postoperative cognitive dysfunction (POCD); hyperglycemia; delirium; neurocognitive; Cerebrospinal fluid (CSF); sorbitol; fructose; thoracic aortic surgery; Montreal Cognitive Assessment (MoCA)

Liquor cerebrospinalis; hersenvocht; polyol route; cerebraal glucose metabolisme; hyperglykemie; delier; postoperatieve cognitieve dysfunctie (POCD); thoracale aorta chirurgie

Ondersteuning

Primaire sponsor: Sponsor: Amsterdam UMC. location AMC, Department of Anesthesiology

Overige ondersteuning: Subsiding party: European Society of Anesthesiology (ESA)

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The main outcome measures per aim are:

- The changes in cerebral glucose metabolism in the perioperative period as measured by the difference in CSF/plasma ratio* of glucose at different points in time

- The increase in cerebral sorbitol and fructose concentrations in the perioperative period as measured by the difference in CSF/plasma ratio of sorbitol and fructose at different points in time

*The CSF/plasma glucose ratio is used because of our hypothesis that an immediate reduction in cerebral hexokinase activity will cause an increase in cerebral glucose and therefore an increase in the CSF/plasma glucose ratio.

Toelichting onderzoek

Achtergrond van het onderzoek

Along with the intended depression of the state of consciousness, anaesthetic drugs decrease cerebral glucose metabolism by 25-63%. Paradoxically, plasma glucose increases, because of the surgical stress response and accompanied insulin resistance. This will lead to an acute supply-demand mismatch and excess of glucose in the brain, and we hypothesize that this leads to activation of the neurotoxic polyol pathway. Postoperative cognitive dysfunction (POCD) has been reported in relation to hypoglycemia. POCD is also suggested to be associated with hyperglycemia, although this is less constantly reported. This hypothesized activation of the neurotoxic polyol pathway may contribute to the relation between hyperglycemia during anaesthesia and postoperative cognitive dysfunction, delirium, and other neurocognitive complications.

Doel van het onderzoek

As plasma glucose during surgery increases, because of the surgical stress response and accompanied insulin resistance, this will lead to an acute supply-demand mismatch and excess of glucose in the brain. We hypothesize that this leads to activation of the neurotoxic polyol pathway. This hypothesized activation of the neurotoxic polyol pathway may contribute to the relation between hyperglycemia during anaesthesia and postoperative cognitive dysfunction, delirium, and other neurocognitive complications.

Onderzoeksopzet

A fasting preoperative plasma glucose is determined.

We will collect the first CSF sample from the reservoir 15 minutes before induction of anaes-

thetia ($t=-15$ min), when the patient has had an overnight fast.

When a CSF sample is obtained, a paired plasma sample will be collected from the arterial line.

The second samples are collected before the start of cardiopulmonary bypass and the third samples will be collected after stopping cardiopulmonary bypass. After discharge from the operating theatre, CSF and plasma samples will be taken every morning before breakfast (i.e. fasting) until removal of the catheter, at latest 48 hours after surgery.

An estimated 4-5 repeated paired measurements per patient would be collected.

We will administer a questionnaire to assess pre- and postoperative cognitive dysfunction, as measured by the Montreal Cognitive Assessment (MoCA). The questionnaire is administered on the day before surgery at approximately 1pm and two weeks after surgery at approximately 1pm.

Onderzoeksproduct en/of interventie

No interventions.

Contactpersonen

Publiek

J. Hermanides
[default]
The Netherlands
0031-205669111

Wetenschappelijk

J. Hermanides
[default]
The Netherlands
0031-205669111

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Patients undergoing elective thoracic aortic surgery without pre-existing brain disease and without diabetes mellitus who are able and willing to participate in the study and can provide written informed consent. Patients must be 18 years or older.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Patients undergoing elective thoracic aortic surgery with pre-existing brain disease and/or diabetes mellitus. Patients who are unable to understand or fill-in questionnaires in Dutch will also be excluded from the study.

Onderzoeksopzet

Opzet

Type:	Observationeel onderzoek, zonder invasieve metingen
Onderzoeksmodel:	Anders
Blinding:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-01-2019
Aantal proefpersonen:	16
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	13-12-2018
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 48939

Bron: ToetsingOnline

Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL7427
NTR-old	NTR7669
CCMO	NL65815.018.18
OMON	NL-OMON48939

Resultaten