

# International study in multiple centers studying the effect of hydroxy-urea and temozolomide chemotherapy in patients with a recurrent malignant brain tumor (Glioblastoma)

Gepubliceerd: 14-04-2016 Laatste bijgewerkt: 13-12-2022

Determining maximal tolerated dose hydroxy-ureum in combination with dose intense temozolomide in patients with recurrent glioblastoma

<b>Ethische beoordeling</b>	Niet van toepassing
<b>Status</b>	Werving nog niet gestart
<b>Type aandoening</b>	-
<b>Onderzoekstype</b>	Interventie onderzoek

## Samenvatting

### ID

NL-OMON20720

### Bron

NTR

### Verkorte titel

HUTMZ

### Aandoening

Brain tumor, glioma, glioblastoma, recurrent glioblastoma, glioblastoma multiforme

Hersentumor, glioom, glioblastoom, recidief glioblastoom, glioblastoma multiforme

## Ondersteuning

**Primaire sponsor:** VU University Medical Center

**Overige ondersteuning:** Janivo Stichting  
CCA fonds

# Onderzoeksproduct en/of interventie

## Uitkomstmaten

### Primaire uitkomstmaten

To determine the maximum tolerated dose (MTD) and safety profile of daily hydroxyurea in combination with continuous dose-intense temozolomide in patients with recurrent GBM.

## Toelichting onderzoek

### Achtergrond van het onderzoek

Glioblastoma is the most frequent and most malignant subtype of primary brain tumors. Unfortunately, over the last two decades, the only major development in the treatment of patients with GBM has been the addition of the DNA alkylating agent TMZ to the standard of care including surgery and radiation resulting in an improvement of the overall survival from 12.1 to 14.6 months

1. More than 90% of patients though die within 5 years of diagnosis. This colossal failure has been partially attributed to development of drug resistance. A major predictor of GBM-response to TMZ is the intrinsic MGMT (O6-methylguanine methyl transferase) promoter methylation status. TMZ induces methylation of guanine at O6-position, a change that causes a futile cycle of attempted DNA repair and results in cell apoptosis. MGMT can remove DNA adducts caused by alkylating agents, resulting in resistance to TMZ. Methylation of the MGMT promotor (occurring in approximately half of GBM) leads to a transcriptional silencing of the MGMT-gene. Patients with MGMT methylated promoters are more likely to benefit from the addition of TMZ. Given that all GBM eventually acquire resistance to TMZ, studies are underway to unravel the molecular changes that occur during treatment and to characterize the mechanisms of resistant recurrences. Through drug screening, we identified HU to sensitize TMZ-resistant GBM cells to TMZ, both in culture and in primary GBM in vivo intracranial models.

We also explored the effect of mechanism behind the synergistic effect of HU+TMZ. HU acts specifically on the S-phase of the cell cycle by inhibiting the enzyme ribonucleotide reductase (RNR), thereby hindering the reductive conversion of ribonucleotides to deoxyribonucleotides and thus limiting de novo DNA synthesis. This property makes RNR inhibitors like HU an attractive candidate for cancer therapy. Another advantage for the use of HU in GBM is that it increases the blood brain tumor (BBB) permeability of certain chemotherapeutics and therefore could enhance penetration of TMZ into GBM. The Notch pathway is another suggested target for cancer therapy and Notch inhibitors have been reported to synergize with TMZ in GBM. In preliminary experiments we observed that HU also was able to inhibit Notch signaling. We are currently, trying to unravel the exact mechanism through which HU achieves re-sensitization for TMZ. Through its inhibitory effects on the Notch pathway and RNR and possibly increasing permeability of the BBB to TMZ, HU has effects on multiple

levels to sensitize cells to TMZ. Our next step is to investigate the safety and MTD of the combination HU+TMZ in patients with rGBM.

1 R Stupp et al. Radiotherapy plus Concomitant and Adjuvant Temozolomide for Glioblastoma. N Engl J Med 2005; 352:987-96;

2 ME Hegi et al. MGMT Gene Silencing and Benefit from Temozolomide in Glioblastoma. N Engl J Med 2005; 352:997-1003;

3 JR Perry et al. Phase II Trial of Continuous Dose-Intense Temozolomide in Recurrent Malignant Glioma: RESCUE Study. JCO 28;12::2051-7 ;

4 C Hunter et al. A hypermutation phenotype and somatic MSH6 mutations in recurrent human malignant gliomas after alkylator chemotherapy. Cancer Res. 2006;66:3987-91.

### **Doel van het onderzoek**

Determining maximal tolerated dose hydroxy-ureum in combination with dose intense temozolomide in patients with recurrent glioblastoma

### **Onderzoeksopzet**

Start 01-06-2016, end 01-06-2020

### **Onderzoeksproduct en/of interventie**

Combine hydroxy-ureum with dose-dense temozolomide in patients with recurrent glioblastoma.

## **Contactpersonen**

### **Publiek**

[default]

The Netherlands

### **Wetenschappelijk**

[default]

The Netherlands

# Deelname eisen

## Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Participants must have histologically or cytologically confirmed glioblastoma multiforme
2. Patients may have had any number of prior therapies for glioblastoma. Patients must be at least 28 days from any investigational agent, 28 days from prior cytotoxic therapy (except 23 days from prior temozolomide, 14 days from vincristine, 42 days from nitrosoureas, 21 days from procarbazine administration), and 7 days for patients who received metronomic chemotherapy or non-cytotoxic agents, e.g., interferon, tamoxifen, thalidomide, cis-retinoic acid, etc.
3. Age  $\geq 18$  years. Because no dosing or adverse event data are currently available on the use of hydroxyurea in combination with temozolomide in participants  $<18$  years of age, children are excluded from this study, but will be eligible for future pediatric trials.
4. Karnofsky Performance Status (KPS)  $\geq 60\%$
5. Participants must have normal organ and marrow function as defined below:
  - leukocytes  $\geq 3,000/\text{mCL}$
  - absolute neutrophil count  $\geq 1,500/\text{mCL}$
  - platelets  $\geq 100,000/\text{mCL}$
  - total bilirubin within normal institutional limits
  - AST(SGOT)/ALT(SGPT)  $\leq 2.5 \times$  institutional upper limit of normal
  - creatinine below upper limit of normal institutional limitsOR
  - creatinine clearance  $\geq 60 \text{ mL/min/1.73 m}^2$  for patients with creatinine levels above institutional normal.
6. Progressive disease on contrast-enhanced brain CT or MRI as defined by RANO Criteria [22], or have documented recurrent glioblastoma on diagnostic biopsy. Patients who have been previously treated with bevacizumab therapy that have T2-weighted or FLAIR MRI sequences considered to be progressive disease by the study investigator but have no contrast-enhancing areas of recurrent disease are eligible.

7. Interval of at least 2 weeks from any prior neurosurgical resection (1 week for intracranial biopsy) to start of study drug; and patient must have adequate wound healing.
8. Interval of at least 12 weeks from prior radiotherapy unless there is either: a) histopathologic confirmation of recurrent tumor, or b) new enhancement on MRI outside of the radiation treatment (RT) field.
9. Because cytotoxic agents such as temozolomide and hydroxyurea are known to be teratogenic, women of child-bearing potential and men must agree to use adequate contraception (hormonal or barrier method of birth control; abstinence) prior to study entry and for the duration of study participation. Should a woman become pregnant or suspect she is pregnant while she or her partner is participating in this study, she should inform her treating physician immediately. Men treated or enrolled on this protocol must also agree to use adequate contraception prior to the study, for the duration of study participation, and 4 months after completion of temozolomide and hydroxyurea administration.
10. Patients must have archival tumor tissue available for molecular analysis and must be willing to consent for this tissue to be analyzed as part of this study. However, if no archival tumor tissue is available, the patient will not be excluded from the study.
11. Ability to understand and the willingness to sign a written informed consent document.

### **Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)**

1. Participants who are receiving any other investigational agents or devices in investigation for glioblastoma.
2. Patients must not have been previously treated with an anti-VEGF inhibitor.
3. History of allergic reactions attributed to compounds of similar chemical composition to temozolomide and/or hydroxyurea.
4. Uncontrolled intercurrent illness including, but not limited to, ongoing or active infection, symptomatic congestive heart failure, unstable angina pectoris, cardiac arrhythmia, or psychiatric illness/social situations that would limit compliance with study requirements.
5. Pregnant women are excluded from this study because hydroxyurea and temozolomide have known teratogenic or abortifacient effects. Because there is an unknown but potential risk for adverse events in nursing infants secondary to treatment of the mother with hydroxyurea and temozolomide, breastfeeding should be discontinued if the mother is treated with hydroxyurea and temozolomide.
6. HIV-positive participants on combination antiretroviral therapy are ineligible because of the potential for pharmacokinetic interactions with hydroxyurea. In addition, these participants

are at increased risk of lethal infections when treated with marrow-suppressive therapy. Appropriate studies will be undertaken in participants receiving combination antiretroviral therapy when indicated.

7. Patients with a history of a different malignancy are ineligible except for the following circumstances: if they have been disease-free for at least 3 years and are deemed by the investigator to be at low risk for recurrence of that malignancy; patients with treated cervical cancer in situ, and basal cell or squamous cell carcinoma of the skin. Patients will not be eligible if they have evidence of other malignancy requiring therapy other than surgery within the last 3 years.

8. Major surgery (not including minor diagnostic procedures such as lymph node biopsy) within 2 weeks of start of study drug; or not fully recovered from any side effects of previous procedures.

9. Presence of extra-cranial metastatic disease.

## Onderzoeksopzet

### Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Anders
Toewijzing:	N.v.t. / één studie arm
Blinding:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

### Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-06-2016
Aantal proefpersonen:	35
Type:	Verwachte startdatum

## Ethische beoordeling

Niet van toepassing

Soort: Niet van toepassing

## Registraties

### Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

### Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

### In overige registers

Register	ID
NTR-new	NL5768
NTR-old	NTR6009
Ander register	: Pro-16/11

## Resultaten