

In vivo study of pulse conductance of oxygenators during pulsatile flow

Gepubliceerd: 17-06-2008 Laatste bijgewerkt: 13-12-2022

The objective of this clinical study is to endorse the hypothesis that oxygenators with a relatively high compliance and relatively low hydraulic resistance enable better pulse conductance than relatively stiff oxygenators featuring relatively high...

Ethische beoordeling	Positief advies
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON20861

Bron

NTR

Verkorte titel

Pulse Conductance of Oxygenators

Aandoening

oxgenator
pulsatile flow
heart-lung machine

Ondersteuning

Primaire sponsor: Jos G. Maessen, MD PhD

University Hospital Maastricht

Dept. Cardiothoracic Surgery

P. Debeyelaan 25

P.O. box 5800

NL - 6200 AZ Maastricht

+31 43 3877070

antoinepsimons@ctc.unimaas.nl

Overige ondersteuning: Jos G. Maessen, MD PhD

University Hospital Maastricht

Dept. Cardiothoracic Surgery
P. Debeyelaan 25
P.O. box 5800
NL - 6200 AZ Maastricht
+31 43 3877070
antoinepsimons@ctc.unimaas.nl

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

To measure pulse conductance of two standard types of oxygenators standard used during coronary artery bypass surgery using a standard heart-lung machine.

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale: The objective of this clinical study is to endorse the hypothesis that oxygenators with a relatively high compliance and relatively low hydraulic resistance enable better pulse conductance than relatively stiff oxygenators featuring relatively high resistance.

Objective: Primary objective: measure pulse conductance of two standard types of oxygenators standard used during coronary artery bypass surgery using a standard heart-lung machine.

Secondary objective: measure/record the standard normalized index of haemolysis during elective coronary artery bypass surgery using a standard heart-lung machine.

Study design: prospective controlled randomized observational study

Study population: 40 patients undergoing elective coronary artery bypass grafting (CABG) using extracorporeal circulation.

Intervention (if applicable): no intervention, standard CABG procedure only

Main study parameters/endpoints: Pulse conductance of the oxygenator used during pulsatile perfusion with a standard heart-lung machine, and the normalized index of haemolysis measured in relation to the oxygenator used during pulsatile perfusion with a standard heart-lung machine.

Nature and extent of the burden and risks associated with participation, benefit and group relatedness: No other or additional risk than for those patients that do not involve in the study, and that undergo coronary artery bypass surgery. Differences in the two groups of standard oxygenators regarding primary and secondary objective is to be investigated. Both devices, however, have been clinically used for more than 15 years, and have been established as equivalent devices to take over lung function during heart surgery using extracorporeal circulation.

Regarding the results of the in vitro investigation in which the Quadrox had a better pulse conductance than the Capiiox SX18, we expect a better pulse conductance in the Quadrox group in the clinical study as well. This would imply a less aggressive pulse generation in the Quadrox group, thus a reduced haemolysis. In standard clinical (non investigational) treatment, however, over the last 15 years, the haemolysis when using the Capiiox SX18 oxygenator has not stood out. Therefore, we do not expect any complications in this group.

Doel van het onderzoek

The objective of this clinical study is to endorse the hypothesis that oxygenators with a relatively high compliance and relatively low hydraulic resistance enable better pulse conductance than relatively stiff oxygenators featuring relatively high resistance.

Onderzoeksopzet

Baseline measurement of free hemoglobin before initiation of cardiopulmonary bypass and after 30 min on cardiopulmonary bypass; continuous recording of line pressures after initiation of cardiopulmonary bypass.

Onderzoeksproduct en/of interventie

Non-interventional study, therefore non-applicable

Contactpersonen

Publiek

University Hospital Maastricht
Dept. Cardiothoracic Surgery
P.O. box 5800
Antoine P. Simons
P. Debeyelaan 25
Maastricht 6200 AZ
The Netherlands
+31 (0)43 3881066

Wetenschappelijk

University Hospital Maastricht
Dept. Cardiothoracic Surgery
P.O. box 5800
Antoine P. Simons
P. Debeyelaan 25
Maastricht 6200 AZ
The Netherlands
+31 (0)43 3881066

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Patients scheduled for elective CABG using a standard heart-lung machine with either a standard used Maquet Quadrox HMO 2000 oxygenator, or a standard clinically equivalent Terumo Capiiox SX18 oxygenator, using standard pulsatile perfusion at the university hospital Maastricht.
2. Calculated bypass flow of 5 l/min.
3. Written informed consent.
4. Age of 18 and older.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. All other patients that do not comply with the inclusion criteria given above, e.g. patients receiving valve surgery, or combined CABG / valve surgery.
2. No written informed consent.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-07-2008
Aantal proefpersonen:	40
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	17-06-2008
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL1298
NTR-old	NTR1346
Ander register	ABR NL22896.068.08 : MEC 08-2-036
ISRCTN	ISRCTN wordt niet meer aangevraagd

Resultaten