

Lifestyle Interventions for Severe Mentally Ill Outpatients in the Netherlands.

Gepubliceerd: 21-12-2012 Laatste bijgewerkt: 13-12-2022

1. A multidimensional lifestyle intervention, including aspects of increased awareness, motivation, self-management, diet, exercise, and a supportive environment, improves or prevents deterioration of the cardiometabolic risk factors of SMI...

Ethische beoordeling	Positief advies
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON21055

Bron

NTR

Verkorte titel

LION

Aandoening

Cardiometabolic health of severe mentally ill outpatients

Ondersteuning

Primaire sponsor: GGZ Friesland, University Medical Center Groningen

Overige ondersteuning: ZonMW

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Waist circumference.

Toelichting onderzoek

Achtergrond van het onderzoek

Patients with severe mental illness (SMI) have an increased cardiometabolic risk. The prevalence of obesity is 4-5 times higher than in the general population, between 45-55%, and the prevalence of type diabetes is around 10-15%, which is up to four times higher than in the general population of comparable age. Although monitoring the somatic health of SMI patients is now obligatory in The Netherlands, most comorbidities are left untreated due to a lack of knowledge, and fear that medication will interact with antipsychotic medication. Lifestyle intervention in high risk individuals from the general population has been shown to be (cost-)effective, and even more effective than early pharmacological treatment, to prevent type 2 diabetes and reduce cardiometabolic risk. This non-pharmacological intervention to reduce cardiometabolic risk may also be effective in SMI patients. In spite of the much increased attention for their excessive cardiovascular risk, evidence based strategies that can be implemented on a large scale to prevent the burden of somatic disease in severe mentally ill (SMI) patients are still lacking, partly because RCTs have been small and of short duration, or not feasible for large-scale implementation. The aim of the current proposal is to compare the (cost)effectiveness of a 12-month multidimensional lifestyle approach for SMI outpatients to usual care to reduce cardiometabolic risk factors in SMI patients. The intervention is based on a state-of-the-art intervention design, promoting active self-management using e-health tools and incorporating motivational techniques by nurses on top of exercise and other health promotion sessions and support from nurses. Secondary research questions include whether the intervention decreases depressive and negative symptoms and whether the multidimensional lifestyle approach is cost-effective. The self-management tools based on e-health are the Traffic Light website, and Heartville, a serious health game that was awarded the "Game for Health" on the TEDx 2012 in Maastricht, The Netherlands. Several steps guide patient and nurse to better lifestyle habits. First, the Traffic Light method displays a risk profile with all lifestyle behaviors in green, orange or red, depending on the level of risk. The website-generated lifestyle profile provides insight for patient and nurse. Second, the patient decides which behavior he/she wants to change. The nurses use motivational interviewing (MI) techniques and the stages of change model to assist the patient in this process. Third, the Traffic Light is used to create a lifestyle plan, in which patients set their own goals on the chosen lifestyle areas, including what the patient's needs are to achieve the goals, such as family involvement. The nurse's role is to support patients in setting realistic goals. Finally, the Traffic Light model is used to sustain change: nurses support patients in the various phases of trial and error, and nurses will use MI and Stages of Change techniques at every step. In addition, the Traffic Light method contains features to support the role of the nurse in ensuring the

availability of payable exercise and health promotion activities, and up to date lifestyle knowledge in the team.

In this multicenter study, it is foreseen that 16 teams from 5 locations will deliver 32 nurses and 640 SMI outpatients. It is a cluster randomized trial, since teams will be randomized, and data analysis will be clustered. The primary outcome measure is waist circumference. Secondary outcomes include readiness to change (motivation), weight and height to calculate BMI, and other risk factors including all the components of the metabolic syndrome. These include blood pressure, plasma triglycerides, cholesterol (LDL, HDL, and total cholesterol) and fasting glucose and HbA1c. Furthermore, the measurements include depressive symptoms, negative symptoms, quality of life, and registration of antipsychotic medication. These measures are part of standard Routine Outcome Monitoring measurements and are taken at baseline, after 6 and after 12 months. Additional assessment of weight, waist circumference and lifestyle is taken at 3 and 9 months for intervention participants to evaluate lifestyle progress. Cost-effectiveness analysis (CEA) takes into account care consumption, waist circumference and quality of life. A budget impact analysis is performed by extrapolating CEA results to yearly intervention costs.

In the period between obtaining funding and preparing the study, unexpected large changes in the organisation of mental health care took place. Budgets were restrained and care delivery shifted towards general mental health care, leading to necessary adjustments in study design. The initial sample size was estimated based on 64 nurses all including 10 patients leading to an target sample of 640 patients. The Medical Ethics Committee advised us to plan an extra 20% inclusion to account for clustering of the data, yielding a target sample of 768 patients. However, it became clear that inclusion of twelve patients per nurse was too high for proper implementation of the intervention due to increased workload of nurses. To compensate, more nurses were trained in order to include less patients per nurse, and inclusion criteria were broadened so that patients in sheltered living facilities could be included as well. With a sample of 250 patients, we are able to detect a significant reduction of 5.8% in waist circumference (primary outcome) and a reduction of 0.6 mmol/L in plasma glucose ($\alpha = 0.05$, power 0.80), leading to a target sample of 250 patients. To correct for drop-out we aimed to include an additional 10%. The funding agency (ZonMW) and the Medical Ethics Committee have approved the adjusted study design and adjusted final sample size of 275 patients.

Doel van het onderzoek

1. A multidimensional lifestyle intervention, including aspects of increased awareness, motivation, self-management, diet, exercise, and a supportive environment, improves or prevents deterioration of the cardiometabolic risk factors of SMI outpatients;
2. The intervention decreases depressive and negative symptoms;
3. The multidimensional lifestyle approach is cost-effective.

Onderzoeksopzet

Measurement before start of trial (T0), after 6 months (T1) and after 12 months, end of trial (T2).

Onderzoeksproduct en/of interventie

The intervention is based on a state-of-the-art intervention design, promoting active self-management using e-health tools and incorporating motivational techniques by

nurses on top of exercise and other health promotion sessions and support from nurses.

The self-management tools based on e-health are the Traffic Light website, and Heartville, a serious health game that was awarded the “Game for Health” on the TEDx 2012 in Maastricht, The Netherlands. Several steps guide patient and nurse to better lifestyle habits. First, the Traffic Light method displays a risk profile with all lifestyle behaviors in green, orange or red, depending on the level of risk. The website-generated lifestyle profile provides insight for patient and nurse. Second, the patient decides which behavior he/she wants to change. The nurses use motivational interviewing (MI) techniques and the stages of change model to assist the patient in this process. Third, the Traffic Light is used to create a lifestyle plan, in which patients set their own goals on the chosen lifestyle areas, including what the patient’s needs are to achieve the goals, such as family involvement. The nurse’s role is to support patients in setting realistic goals. Finally, the Traffic Light model is used to sustain change: nurses support patients in the various phases of trial and error, and nurses will use MI and Stages of Change

techniques at every step. In addition, the Traffic Light method contains features to support the role of the nurse in ensuring the availability of payable exercise and health promotion activities, and up to date lifestyle knowledge in the team.

The intervention will take one year.

The intervention will be performed by nurses of FACT teams, who are specially trained. Every institution has at least 2 FACT teams. These teams will be randomised into training (two nurses per team) or no training (control condition).

Contactpersonen

Publiek

GGZ Friesland
Postbus 932
Frederike Jörg
Leeuwarden 8901 BS
The Netherlands
+31 (0)58 2848985

Wetenschappelijk

GGZ Friesland
Postbus 932
Frederike Jörg
Leeuwarden 8901 BS
The Netherlands
+31 (0)58 2848985

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Severe mentally ill outpatients with at least one of the following risk factors:

1. Waist circumference > 102 cm (m) or > 88 cm (f);
2. Fasting glucose > 5.6 mmol/l;
3. BMI > 25 kg/m².

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Pregnancy.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blindering:	Enkelblind
Controle:	Geneesmiddel

Deelname

Nederland	
Status:	Werving gestopt
(Verwachte) startdatum:	01-01-2014
Aantal proefpersonen:	275
Type:	Werkelijke startdatum

Voornemen beschikbaar stellen Individuele Patiënten Data (IPD)

Wordt de data na het onderzoek gedeeld: Nog niet bepaald

Ethische beoordeling

Positief advies	
Datum:	21-12-2012
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL3606
NTR-old	NTR3765
Ander register	ZonMw : 837001006
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/Q