Efficacy of peripherally targeted inhaled rhDNase for persistent obstructive asthma in childhood.

Gepubliceerd: 02-05-2006 Laatst bijgewerkt: 13-12-2022

rhDNase improves lung function in children with persistent asthma who have persistent obstructive pulmonary function; Treatment of mucus impaction is an interesting alternative approach to treat peripheral airflow limitation in asthmatic patients....

Ethische beoordeling	Niet van toepassing
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON21320

Bron NTR

Verkorte titel

Aandoening

asthma with persistent obstructive pulmonary function.

Ondersteuning

Primaire sponsor: ErasmusMC Rotterdam **Overige ondersteuning:** E.M. Bakker receives a 4-year grant of Roche Nederland B.V. for her PhD research.

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

1 - Efficacy of peripherally targeted inhaled rhDNase for persistent obstructive ast ... 4-05-2025

Primary endpoint will be the change in FEF75 as a result of treatment. FEF75 is the most suitable endpoint since it is sensitive to peripheral airways obstruction.

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale: One of the aims of the pharmacological management of asthma is to normalize pulmonary function. In fact, in many children with asthma treated with ICS and inhaled b2-agonists, pulmonary function tests return to (nearly) normal values. However, airflow limitation (airways obstruction) persists in a proportion of these patients. The pathogenesis of this persistent obstruction is unclear. Airway wall inflammation and edema, increased bronchomotor tone, increased sputum volume, increased sputum viscoelasticity and decreased mucociliary clearance all play an important role in the pathogenesis of airways obstruction in childhood asthma. Since absolute airway resistance is higher in childhood than in adulthood, it is likely that mucus retention contributes substantially to airways obstruction in childhood asthma.

In severe asthma dramatic improvement has been described in a few patients after inhalation of the mucolytic rhDNase. In addition in pathology studies extensive mucus plugging has been described in asthma patients. Based on these findings we hypothesize that additional treatment benefit can be obtained when mucus plugging is targeted as part of asthma treatment in children with asthma and persistent airways obstruction.

Therefore we would like to investigate the additional effect of rhDNase treatment in children with asthma and persistent obstructive pulmonary function.

Objective: Primary Objective: To investigate the effect of treatment with nebulized rhDNase on pulmonary function in children with asthma and persistent obstructive pulmonary function.

Secondary Objectives: To investigate the effect of treatment with nebulized rhDNase on gas mixing, FENO values and symptom scores in children with asthma and persistent obstructive pulmonary function.

Study design: This study will be a randomized placebo-controlled cross-over clinical trial. Study population: Children with asthma and persistent airway obstruction, aged 6-18 years. Intervention: All patients will be treated with 2 weeks of placebo and 2 weeks of rhDNase, separated by a 4-week washout period. Children will be randomized to start with rhDNase or placebo.

Doel van het onderzoek

rhDNase improves lung function in children with persistent asthma who have persistent obstructive pulmonary function;

Treatment of mucus impaction is an interesting alternative approach to treat peripheral airflow limitation in asthmatic patients. In severe asthma dramatic improvement has been described in a few patients after inhalation of the mucolytic rhDNase. In addition in pathology

studies extensive mucus plugging has been described in asthmatic patients. Based on these findings we think that additional treatment benefit can be obtained when mucus plugging is targeted as part of asthma treatment. Especially for those asthmatic children with persistent peripheral airways obstruction, rhDNase is a well known and safe drug that could be used for this treatment. Therefore we hypothesize that rhDNase has additional effect on lung function in children with persistent asthma who have persistent obstructive pulmonary function.

Onderzoeksproduct en/of interventie

Nebulization with rhDNase or placebo once daily (each participant is treated for 2 weeks with rhDNase and for 2 weeks with placebo).

Contactpersonen

Publiek

Erasmus Medical Center, Sophia Children's Hospital Rotterdam, Department of Pediatric Pulmonology, Dr. Molewaterplein 60 H.A.W.M. Tiddens Dr. Molewaterplein 60 Rotterdam 3015 GJ The Netherlands +31 (0)10 4636690 / +31 (0)10 4636363 (general)

Wetenschappelijk

Erasmus Medical Center, Sophia Children's Hospital Rotterdam, Department of Pediatric Pulmonology, Dr. Molewaterplein 60 H.A.W.M. Tiddens Dr. Molewaterplein 60 Rotterdam 3015 GJ The Netherlands +31 (0)10 4636690 / +31 (0)10 4636363 (general)

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen

(Inclusiecriteria)

1. Age 6 – 18 years;

2. Asthma diagnosed according to GINA guidelines;

3 Attending the outpatient clinic for at least one year;

4.Treatment with at least 400 mg/day inhaled Budesonide or equivalent (dose constant for at least 6 months) and bronchodilators as needed or daily;

5. Clinically stable asthma while using a constant dose of ICS for at least three months;

6. Ability to perform lung function tests (assessed by trained lung function technician);

7. Persistent peripheral airways obstruction as assessed by pulmonary function testing, defined as: dissociation between FVC and FEF75 values: FEF75 at least 20% (absolute % predicted) lower than FVC;

8. FVC within normal limits (for this study defined as FVC > 80% pred).

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- 1. Asthma exacerbation with hospital admission in last 3 months;
- 2. Intensive Care Unit (ICU) admission for asthma within the last year;
- 3. Current respiratory tract infection;
- 4. Inability to follow instructions of the investigator;
- 5. Inability to inhale rhDNase;

6. Concomitant medical conditions that effect inhaled treatment (e.g. cleft palate, severe malacia);

- 7. Neuromuscular disease;
- 8. Smoking.

4 - Efficacy of peripherally targeted inhaled rhDNase for persistent obstructive ast ... 4-05-2025

Onderzoeksopzet

Opzet

Туре:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blindering:	Dubbelblind
Controle:	Placebo

Deelname

Nederland	
Status:	Werving gestopt
(Verwachte) startdatum:	01-09-2006
Aantal proefpersonen:	60
Туре:	Werkelijke startdatum

Ethische beoordeling

Niet van toepassing	
Soort:	Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL612
NTR-old	NTR671
Ander register	: 2412325-2
ISRCTN	ISRCTN71537084

Resultaten

Samenvatting resultaten

1. Fuchs, H.J., et al., Effect of aerosolized recombinant human DNase on exacerbations of respiratory symptoms and on pulmonary function in patients with cystic fibrosis. The Pulmozyme Study Group. N Engl J Med, 1994. 331(10): p. 637-42.

2. King, M., et al., Rheology of cystic fibrosis sputum after in vitro treatment with hypertonic saline alone and in combination with recombinant human deoxyribonuclease I. Am J Respir Crit Care Med, 1997. 156(1): p. 173-7.

3. Hodson, M.E. and P.L. Shah, DNase trials in cystic fibrosis. Eur Respir J, 1995. 8(10): p. 1786-91.

4. Greally, P., Human recombinant DNase for mucus plugging in status asthmaticus. Lancet, 1995. 346(8987): p. 1423-4

5. Durward, A., V. Forte, and S.D. Shemie, Resolution of mucus plugging and atelectasis after intratracheal rhDNase therapy in a mechanically ventilated child with refractory status asthmaticus. Crit Care Med, 2000. 28(2): p. 560-2.