

Effect of occlusal loading on secondary caries development in situ'

Gepubliceerd: 10-09-2019 Laatst bijgewerkt: 13-12-2022

Compressive occlusal loading of a space between composite and tooth material will lead to an increase in lesion depth and mineral loss in these samples.

Ethische beoordeling	Niet van toepassing
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON21363

Bron

Nationaal Trial Register

Verkorte titel

ELCIS

Aandoening

secondary caries

Ondersteuning

Primaire sponsor: Radboudumc

Overige ondersteuning: Stichting Bevordering Tandheelkundige Kennis

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Lesion depth, measured through transversal microradiography

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale:

Secondary caries is tooth decay next to an existing restoration and one of the main reasons to replace restoration in general dental practice. Previous research has shown that secondary caries lesions can develop when a small gap is present between the tooth and restorative material. Gaps as small as 30 μm have been shown to develop secondary caries lesions in vitro and in situ [Kuper et al., 2014; Maske et al., 2019; Maske et al., 2017].

Presence of a gap allows the formation of a wall and surface lesion next to a dental restoration. An in vitro study by Kuper et al. [2013] showed that an increased hydrodynamic flow in the gap between composite and tooth material leads to an increase in wall lesion development.

A more recent study by Askar et al. [2017] found that loading a tooth-restoration sample containing a gap of 100 μm width led to increased caries lesion formation compared to unloaded samples. They theorized that the loading on the sample containing a gap could lead to compression of the gap area. When the force is released, the compression ends, and the gap returns to normal. This cyclic deformation of the gap area could possibly lead to an increased hydrodynamic flow, which explains the increase in lesion formation in loaded samples.

Since only in vitro data is available on this topic, the clinical relevance is so far unclear. Whether the effect of loading of restorations still influences the secondary caries process in a more variable clinical environment, is unknown.

Objective:

The aim of this in situ study is to investigate whether occlusal loading leads to increased secondary caries formation in loaded compared to unloaded samples in a near-clinical environment.

Study design:

A mono-center, single blind in situ study, with split-mouth design.

Edentulous subjects will receive a duplicate of their lower denture containing sterilized tooth samples in slots. Half the samples will be placed in occlusal contact with the upper denture. The other half will be placed slightly lower, out of contact. The appliance has to be dipped in 20% sucrose solution 4 times a day for 10 minutes. The appliance containing samples needs to be worn for 6 weeks, 24 hours a day.

Study population:

Healthy edentulous volunteers (>18 years old) who wear a full denture in the lower jaw. A sample size of $n=14$ will be aimed for.

Main study parameters/endpoints:

The main study parameters are lesion depth (μm) and integrated mineral loss ($\mu\text{m.vol}\%$) in the tooth sample close to the restoration margin. These will be measured through the microradiographic technique T-WIM.

Doel van het onderzoek

Compressive occlusal loading of a space between composite and tooth material will lead to an increase in lesion depth and mineral loss in these samples.

Onderzoeksopzet

Start of wearing denture containing samples = t0

Samples are worn for 6 weeks. Samples are measured afterwards = t42

Onderzoeksproduct en/of interventie

Lower dental prosthesis is copied and equipped with samples. No direct intervention on subjects. Samples are dipped in sucrose solution 4 times a day.

Contactpersonen

Publiek

Radboudumc
Audrey Hollanders

0243618824

Wetenschappelijk

Radboudumc
Audrey Hollanders

0243618824

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

- Adults (>18 years)
- Wear a full denture in the lower jaw
- Have a lower denture height of at least 8 mm

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- ASA score >2
- Unable to give informed consent
- Unable to understand written patient information

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Factorieel
Toewijzing:	N.v.t. / één studie arm
Blinding:	Enkelblind
Controle:	Placebo

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-01-2020
Aantal proefpersonen:	14
Type:	Verwachte startdatum

Voornemen beschikbaar stellen Individuele Patiënten Data (IPD)

Wordt de data na het onderzoek gedeeld: Nee

Ethische beoordeling

Niet van toepassing	
Soort:	Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL8031
Ander register	ABR : 71551

Resultaten