

Het behandelen van kamerritmestoornissen bij patiënten met Tetralogie van Fallot, die een heroperatie ondergaan waarbij de pulmonalisklep wordt vervangen en die risicolopen op kamerritmestoornissen.

Gepubliceerd: 07-10-2013 Laatste bijgewerkt: 19-03-2025

Improved understanding of slow conducting anatomical isthmuses in Tetralogy of Fallot and cryoablation of these slow conducting anatomical isthmuses during pulmonary valve replacement might prevent VT re- and occurrence.

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON21437

Bron

Nationaal Trial Register

Verkorte titel

N/A

Aandoening

Congenital heart disease

Ondersteuning

Primaire sponsor: Leiden University Medical Center (LUMC)

Overige ondersteuning: Leiden University Medical Center (LUMC)

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The main study endpoints are: (1) achievement of bidirectional conduction block after intraoperative cryoablation, (2) reinducibility of isthmus dependent monomorphic VT after intraoperative cryoablation and (3) occurrence and recurrence of ventricular arrhythmias during follow-up.

Toelichting onderzoek

Achtergrond van het onderzoek

Background of the study:

Tetralogy of Fallot (TOF) is the most common severe congenital heart disease and is associated with late morbidity and mortality due to ventricular arrhythmias (VA). Patients with documented or suspected VA usually receive implantable cardioverter defibrillators (ICDs). However, not all VA are life-threatening although an important source of morbidity. In addition, ICDs do not prevent VA therefore additional and/or alternative treatment options are required. Of importance, the majority of VA associated with TOF are monomorphic ventricular tachycardias (VT). We recently could demonstrate that the substrate for the majority of these monomorphic VTs are slow conducting anatomical isthmuses bordered by unexcitable tissue. These slow conducting isthmuses may be the consequence of the initial repair in childhood but may also be due to the abnormal myocardium of the malformation itself. Targeting these isthmuses by catheter ablation has been shown to prevent VT recurrence and is the accepted current approach in clinical practice. Patients after initial total repair of TOF may require a reoperation for pulmonary valve regurgitation. However, simply replacing the valve does not affect the risk for VT. During reoperation potential slow conducting isthmuses can be ablated with the potential to prevent VT recurrence but also VT occurrence and thereby “curing” the isthmus dependent monomorphic VT provided that isthmus block is achieved. Preventive ablation of the slow conducting isthmuses during surgery becomes particular important if pulmonary valve replacement (PVR) by a homograft is performed. In this case, the homograft may cover important parts of the slow conducting isthmus which makes catheter ablation at a later stage impossible and is the most important reason for ablation failure in patients that present with VT after PVR.

Objective of the study:

- 1) To evaluate the feasibility and the acute effect of intraoperative cryoablation of the slow conducting anatomical isthmuses (endpoint bidirectional conduction block) and on the re-inducibility of monomorphic isthmus dependent VT.
- 2) To study the pathomechanism of slow conduction within these isthmuses by comparing histological and electrophysiological characteristics of biopsies in patients after repair of TOF who undergo reoperation and of patients who undergo first total correction.
- 3) To assess the long-term results of intraoperative cryoablation of the slow conducting anatomical isthmuses on recurrence and occurrence of monomorphic VT.

Study design:

A prospective duo-centre cohort study.

Study population:

Group A: Patients with repaired TOF and accepted for PVR from the age of eight years.

Group B: Patients with TOF who undergo first repair. This group serves only as a control group for the histology of the infundibular muscle which removal is part of the surgical repair.

Intervention:

Intraoperative cryoablation of the slow conducting anatomical isthmuses.

Primary study parameters/outcome of the study:

The main study parameters are: (1) prevalence and characteristics of slow conducting anatomical isthmuses, (2) inducibility of monomorphic isthmus related VT before intraoperative cryoablation, (3) histological and electrophysiological characteristics of the biopsies taken from slow conducting isthmuses (reoperation) and the infundibular muscle which is the most frequent location of a potential slow conducting isthmus later in life (initial repair). The main study endpoints are: (1) achievement of bidirectional conduction block after intraoperative cryoablation, (2) reinducibility of isthmus dependent monomorphic VT after intraoperative cryoablation and (3) occurrence and recurrence of VA during follow-up.

Doel van het onderzoek

Improved understanding of slow conducting anatomical isthmuses in Tetralogy of Fallot and cryoablation of these slow conducting anatomical isthmuses during pulmonary valve replacement might prevent VT re- and occurrence.

Onderzoeksopzet

1. Baseline evaluation (electrophysiologic study with right ventricular mapping and biopsies of slow conducting isthmuses and the infundibular muscle).
2. Follow-up: Primary endpoints.

Onderzoeksproduct en/of interventie

The intervention is cryoablation of slow conducting anatomical isthmuses which (potentially) contain critical parts of the VT re-entry circuit.

The first two study endpoints (i, achievement of bidirectional conduction block after intraoperative cryoablation; ii, re-inducibility of isthmus dependent monomorphic VT after intraoperative cryoablation) will be checked during a post-operative electrophysiologic study with mapping of the right ventricle.

The third study endpoint (occurrence and recurrence of ventricular arrhythmias during follow-up) will be achieved by comparing the study cohort with a historical cohort operated/treated by the same team.

Contactpersonen

Publiek

Department of Cardiology (C5-P)
Leiden University Medical Center
Albinusdreef 2

G.F.L. Kapel
Leiden 2333 ZA
The Netherlands
+31 (0) 71 5265389

Wetenschappelijk

Department of Cardiology (C5-P)
Leiden University Medical Center
Albinusdreef 2

G.F.L. Kapel
Leiden 2333 ZA

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Group A: Patients with repaired Tetralogy of Fallot who undergo reoperation for pulmonary valve replacement.

Group B: Patients with Tetralogy of Fallot who undergo primary repair.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Group A: (1) Age younger than 8 years. (2) Inability to comply with the protocol due to hemodynamic instability.

Both groups: (1) Inability to sign informed consent by the patient or his legal representative.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Niet-gerandomiseerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	15-05-2013
Aantal proefpersonen:	61

Type:

Verwachte startdatum

Ethische beoordeling

Positief advies

Datum:

07-10-2013

Soort:

Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 39860

Bron: ToetsingOnline

Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL4033
NTR-old	NTR4199
CCMO	NL41727.058.12
ISRCTN	ISRCTN wordt niet meer aangevraagd.
OMON	NL-OMON39860

Resultaten

Samenvatting resultaten

N/A