

Mentalization Based Treatment for Psychosis

Gepubliceerd: 19-08-2014 Laatste bijgewerkt: 13-12-2022

Patients who receive treatment as usual (TAU) plus mentalization-based treatment for psychosis (MBT-P) will show a greater improvement in social functioning than patients who receive TAU only. This effect is mediated by an increase of social...

Ethische beoordeling	Positief advies
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON21688

Bron

NTR

Verkorte titel

MBT-P

Aandoening

Nonaffective psychotic disorders:
schizophrenia, schizophreniform or schizoaffective disorder (295.x)
delusional disorder (297.1)
brief psychotic disorder (298.8)
psychotic disorder not otherwise specified (298.9).

Ondersteuning

Primaire sponsor: Maastricht University, the Netherlands

Overige ondersteuning: Rivierduinen Institute for Mental Health Care

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Social Functioning (Social Functioning Scale: average score), Timepoints: 0 months, 18 months, 24 months.

Toelichting onderzoek

Achtergrond van het onderzoek

Many patients with nonaffective psychotic disorders (NAPD) suffer impairments in social cognition and an emotional hyperreactivity to social stress, which often lead to decreased social functioning.

Mentalization Based Treatment for Psychosis (MBT-P) is developed to target impaired social cognition. It is a psychodynamic treatment rooted in attachment and cognitive theory developed for the treatment of NAPD. It is expected to improve social cognition and decrease social stress reactivity which in turn are expected to improve social functioning. MBT-P is further expected to increase quality of life and disease awareness, and to reduce substance abuse, positive, anxious and depressive symptoms.

Doel van het onderzoek

Patients who receive treatment as usual (TAU) plus mentalization-based treatment for psychosis (MBT-P) will show a greater improvement in social functioning than patients who receive TAU only.

This effect is mediated by an increase of social cognition and a decrease of social stress reactivity.

Onderzoeksopzet

0 months: baseline measurement

9 months: halfway treatment measurement

18 months: end of treatment measurement

24 months: follow-up measurement

Onderzoeksproduct en/of interventie

-Mentalization Based Treatment for Psychosis

-Treatment as usual for NAPD: psychiatric consults, supportive therapy, welfare aid, case management.

Contactpersonen

Publiek

GGZ Leiden
J.G. Weijers
Rivierduinen Institute for Mental Health Care
Sandifortdreef 19
Leiden 2333 ZZ
The Netherlands
0031-652582271

Wetenschappelijk

GGZ Leiden
J.G. Weijers
Rivierduinen Institute for Mental Health Care
Sandifortdreef 19
Leiden 2333 ZZ
The Netherlands
0031-652582271

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

- Age 18-55
- Informed consent
- Diagnosis of a nonaffective psychotic disorder
- In treatment for at least a year, but no longer than 10 years

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- Intellectual disabilities and/or illiteracy.
- A lack of mastery of the Dutch language.
- Severe addiction to such an extent that inpatient detoxification is necessary. After detoxification the patient is still eligible for participation in the study.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Enkelblind
Controle:	Geneesmiddel

Deelname

Nederland	
Status:	Werving gestopt
(Verwachte) startdatum:	21-08-2014
Aantal proefpersonen:	90
Type:	Werkelijke startdatum

Voornemen beschikbaar stellen Individuele Patiënten Data (IPD)

Wordt de data na het onderzoek gedeeld: Nog niet bepaald

Ethische beoordeling

Positief advies	
Datum:	19-08-2014
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL4588
NTR-old	NTR4747
Ander register	MEC AZM / MU : 13-3-066.5/ab

Resultaten

Samenvatting resultaten

none yet