

Bimodal success.

Gepubliceerd: 03-04-2013 Laatste bijgewerkt: 19-03-2025

One way of providing binaural cues to unilateral CI recipients is to wear an acoustic hearing aid in the contralateral non-implanted ear, known as bimodal hearing. The benefits of bimodal input (e. g. speech perception in noise, localisation,...

Ethische beoordeling	Positief advies
Status	Anders
Type aandoening	-
Onderzoekstype	Observationeel onderzoek, zonder invasieve metingen

Samenvatting

ID

NL-OMON21746

Bron

Nationaal Trial Register

Aandoening

cochlear implant
CI
cochleair implantaat
hearing aid
hoortoestel
hoorapparaat
bimodal fitting
bimodale aanpassing
bimodal benefit
bimodale winst

Ondersteuning

Primaire sponsor: Maastricht University Medical Centre (MUMC+)

P. Debyelaan 25
6229 HX Maastricht
The Netherlands

+31-43-387 65 43

Overige ondersteuning: Maastricht University Medical Centre (MUMC+)

P. Debyelaan 25
6229 HX Maastricht
The Netherlands

+31-43-387 65 43

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The primary study parameter is the degree of bimodal benefit (=the benefit of wearing a CI in combination with a contralateral hearing aid in comparison to only wearing a unilateral CI). Speech understanding in noise is considered to be the primary outcome measure of bimodal benefit.

In part A this will subjectively be measured using two daily-life disability questionnaires.
In Part B the degree of bimodal benefit will objectively be measured on a spatial speech understanding in noise test.

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale:

Since the 1990s, cochlear implantation (CI) has been the standard practice to restore hearing in severely hearing-impaired and deaf patients by providing multi-channel electrical stimulation to the auditory nerve. Current reimbursement regulations in the Netherlands only allow standard unilateral implantation. However, there is a growing interest in bilateral cochlear implantation and in the benefits of binaural hearing (hearing with two ears) in general. One way of providing binaural cues to unilateral CI recipients is to wear an acoustic hearing aid in the contralateral non-implanted ear, known as bimodal hearing.

The benefits of bimodal input (e. g. speech perception in noise, localisation, listening effort) can objectively be demonstrated in clinical research settings. However, in daily practice there are often patients who choose not to wear their hearing aid (anymore) after receiving a CI in the other ear. Why this is and how this can be improved is still an unanswered but very relevant question in fitting and advising these patients in clinical practice.

Objective:

This research project has 2 main goals and can therefore be divided into 2 parts:

The aim of part (A) is to qualitatively and quantitatively assess the daily-life experiences of CI-patients who do or do not wear a contralateral hearing aid. Assessing the subjective experiences of these patients is a research area that hasn't been covered in literature yet.

The aim of part (B) is to identify the key contributors to successful bimodal fitting. There is still little known about the reasons why patients may or may not experience benefit from wearing a contralateral hearing aid.

After the assessment of these key contributors to bimodal success, a protocol will be developed that may result in a faster and better bimodal fitting. In future research these new insights may lead to clinical tools to create more successful bimodal users with improved hearing performance. Moreover, the results from this study will probably broaden the inclusion criteria for CI in combination with HA, making it available for a larger population of patients.

Study population:

This study focuses on severely hearing impaired adults who use a unilateral cochlear implant with or without a contralateral hearing aid (total n=70).

Study design:

In the first part of this study (A), unilateral CI-users will be asked to fill out a set of relevant daily-life questionnaires in the field of hearing experiences, sound quality and general quality of life. In this way their subjective experiences with wearing or not wearing a hearing aid in the contralateral ear will be reviewed.

In the second part of this study (B), the subset of bimodal users will be asked to partake in objective testing. The aim of this part is to investigate correlations between the benefit of their bimodal fitting (e.g. speech-in-noise and listening effort) and influencing factors (e.g. residual hearing, device fitting, spectral- and temporal resolution and overlap, loudness balance and cognitive skills).

Nature and extent of the burden and risks associated with participation, benefit and group relatedness:

Subjects participating in this study will be asked to fill out a set of questionnaires (A) and/or perform audiological-specific tests (B). Study associated risks are thought of as non-existent

since this is an observational study with non-invasive or low impact tests and no direct interventions. Participation however takes time and effort from subjects.

The aim of the study is to identify key contributors to bimodal success. One may expect that based on these results the clinical practice of unilateral CI patients can be improved, both on an individual and on a group basis. The clinical goal is to enable patients to successfully use CI and HA together, receiving optimal benefit from their residual auditory abilities.

Doel van het onderzoek

One way of providing binaural cues to unilateral CI recipients is to wear an acoustic hearing aid in the contralateral non-implanted ear, known as bimodal hearing.

The benefits of bimodal input (e. g. speech perception in noise, localisation, listening effort) can objectively be demonstrated in clinical research settings. However, in daily practice there are often patients who choose not to wear their hearing aid (anymore) after receiving a CI in the other ear. Why this is and how this can be improved is still an unanswered but very relevant question in fitting and advising these patients in clinical practice.

The hypothesis of this study is that more insight into the subjective experiences of these patients and the factors underlying bimodal benefit will help in developing bimodal fitting guidelines.

Onderzoeksopzet

N/A

Onderzoeksproduct en/of interventie

N/A

Contactpersonen

Publiek

Maastricht UMC+

ENT/Audiological Centre

PO Box 5800
E.M.J. Devocht
Maastricht 6202 AZ
The Netherlands
+31 (0)43 3874594

Wetenschappelijk

Maastricht UMC+

ENT/Audiological Centre

PO Box 5800
E.M.J. Devocht
Maastricht 6202 AZ
The Netherlands
+31 (0)43 3874594

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Capacitated adult (>18 years of age);
2. Patient of CI-team South-East Netherlands;
3. User of a unilateral cochlear implant (CI) of the brand Advanced Bionics;
4. First fit of CI \geq one year ago;
5. Wearing CI speech processor (almost) always (i.e. ≥ 10 hours a day);
6. Agreed to participate in the study (informed consent);
7. PART A: Willing and able to fill out questionnaires;
8. PART B: Wearing a contralateral hearing aid $>50\%$ of the time (≥ 5 hours a day);
9. PART B: Willing and able to visit the azM and participate in testing.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Non Dutch speaking;
2. <18 years of age or incapacitated;
3. Bilateral cochlear implant user (CI+CI);

4. PART B concerning CBCT-scan: Pregnancy, claustrophobia, inability to sit still for longer than 30sec.

Onderzoeksopzet

Opzet

Type:	Observationeel onderzoek, zonder invasieve metingen
Onderzoeksmodel:	Parallel
Toewijzing:	N.v.t. / één studie arm
Blinding:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Anders
(Verwachte) startdatum:	08-04-2013
Aantal proefpersonen:	70
Type:	Onbekend

Ethische beoordeling

Positief advies	
Datum:	03-04-2013
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 39736
Bron: ToetsingOnline
Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL3746
NTR-old	NTR3932
CCMO	NL42011.068.13
ISRCTN	ISRCTN wordt niet meer aangevraagd.
OMON	NL-OMON39736

Resultaten

Samenvatting resultaten

N/A