

Exploration of implicit biases in adults who are current cocaine users

Gepubliceerd: 10-03-2017 Laatste bijgewerkt: 13-12-2022

1. A decrease in cocaine use over a three month period in participants who receive AAT training as opposed to VPT training. 2. A change in symptoms involving craving, the urge to use, withdrawals well as stated readiness to change in participants...

Ethische beoordeling	Niet van toepassing
Status	Werving tijdelijk gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON22059

Bron

Nationaal Trial Register

Verkorte titel

CBM Cocaine SA

Aandoening

Cocaine-dependent disorder as defined in the DSM-5 in South Africa

Ondersteuning

Primaire sponsor: University of Amsterdam (UvA)

Overige ondersteuning: Not applicable

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

A reduction in the number of lines of cocaine used in the past two weeks as measured by a two-week Time Line Feedback (TLFB).

Toelichting onderzoek

Achtergrond van het onderzoek

12-nov-2020: This trial is discontinued.

This research is the first online study targeting cocaine use in South Africa and involving all race groups. Cocaine use is extremely prevalent in urban milieus in South Africa.

Doel van het onderzoek

1. A decrease in cocaine use over a three month period in participants who receive AAT training as opposed to VPT training.
2. A change in symptoms involving craving, the urge to use, withdrawals well as stated readiness to change in participants receiving either AAT or VPT training as opposed to the participants in the placebo group.
3. A correlation between a decrease in cocaine use and the use of CBT coping skills as opposed to meditation-based coping skills.
4. No relationship between Race (Black, White, Mixed race or Colored, Indian) and an outcome defined as reduced cocaine use.

Onderzoeksopzet

Intake, six weeks post-treatment; three months post-treatment.

Onderzoeksproduct en/of interventie

The participants are assessed at intake to determine their level of attentional bias (AB) and approach bias (AAT) to pictorially presented cocaine-related stimuli. Thereafter, they undergo a six-week training component in either attentional bias modification or approach bias modification. They are re-assessed at six weeks and three months post-treatment.

Contactpersonen

Publiek

UvA
Evelyn Beneke
[default]

The Netherlands
021 438 1022 / 082 494 4882

Wetenschappelijk

UvA
Evelyn Beneke
[default]
The Netherlands
021 438 1022 / 082 494 4882

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Self-rated fluency in English; Age 18-65; Willingness to participate

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

History of head injury; color-blindness; photo-sensitive epilepsy

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Dubbelblind
Controle:	Placebo

Deelname

Nederland
Status: Werving tijdelijk gestopt

(Verwachte) startdatum: 04-11-2015
Aantal proefpersonen: 160
Type: Verwachte startdatum

Voornemen beschikbaar stellen Individuele Patiënten Data (IPD)

Wordt de data na het onderzoek gedeeld: Nog niet bepaald

Ethische beoordeling

Niet van toepassing
Soort: Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL6114
NTR-old	NTR6253
Ander register	METC UvA : 2015-DP 4666

Resultaten