

Validation of pelvic floor related symptom and Quality of Life questionnaires in Dutch patients.

Gepubliceerd: 28-05-2010 Laatste bijgewerkt: 18-08-2022

Pelvic floor related symptom and Quality of Life questionnaires (IIQ-7, UDI-6, IIEF-5, PISQ-12, PFIQ-7, PFDI-20, FISQ, FIQL) in Dutch are valid instruments, which can be used reliably in daily practice and clinical research.

Ethische beoordeling	Positief advies
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Observationeel onderzoek, zonder invasieve metingen

Samenvatting

ID

NL-OMON22262

Bron

Nationaal Trial Register

Aandoening

urinary incontinence, erectile dysfunction, pelvic organ prolapse, fecal incontinence

Ondersteuning

Primaire sponsor: Erasmus Medical Center, the Netherlands, department of Urology

Overige ondersteuning: Erasmus Medical Center, the Netherlands, department of Urology

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Questionnaires' Psychometric Properties:

1. Reliability: internal consistency;

2. Reproducibility: test-retest reliability;

3. Validity: convergent validity;

4. Responsiveness: sensitivity of change.

Toelichting onderzoek

Achtergrond van het onderzoek

The objective of this study is to validate the Dutch versions of Pelvic Floor Related Questionnaires (Urogenital Distress Inventory Short Form: UDI-6; Incontinence Impact Questionnaire Short Form: IIQ-7, International Index of Erectile Function short form: IIEF-5, Pelvic Floor Distress Inventory Short Form: PFDI-20, Pelvic Floor Impact Questionnaire Short Form: PFIQ-7, Pelvic Organ Prolapse/Urinary Incontinence Sexual Questionnaire Short Form: PISQ-12, Fecal Incontinence Quality of Life Scale: FIQL, Fecal Incontinence Severity Index: FISI).

The linguistic validation of the questionnaires will be performed through a multistep process: backward and forward translations coordinated by clinical investigators, followed by a pretest. The final versions will be administered to a larger sample of patients, aged 18 years or older, with complaints for at least 3 months. To evaluate test-retest reliability, patients will be re-rated after 1 week. To test the questionnaires' capacity to discriminate patients with or without symptoms (cases and controls, respectively) a sample of 50 healthy patients will be enrolled.

To test for convergent validity, a voiding diary, defecation diary, SF-12 and EuroQol-5 will be used.

To measure the sensitivity of change; questionnaires will be filled out 3 months post therapy.

Hypothesis: Pelvic floor related symptom and Quality of Life questionnaires (IIQ-7, UDI-6, IIEF-5, PISQ-12, PFIQ-7, PFDI-20, FISI, FIQL) in Dutch are valid instruments, which can be used reliably in daily practice and clinical research.

Doel van het onderzoek

Pelvic floor related symptom and Quality of Life questionnaires (IIQ-7, UDI-6, IIEF-5, PISQ-12, PFIQ-7, PFDI-20, FISI, FIQL) in Dutch are valid instruments, which can be used reliably in daily practice and clinical research.

Onderzoeksopzet

1. Baseline;
2. +1 week;
3. Three months after therapy.

Onderzoeksproduct en/of interventie

Dutch version of the questionnaires:

1. Incontinence Impact Questionnaire Short Form (IIQ-7);
2. Urogenital Distress Inventory Short Form (UDI-6);
3. International Index of Erectile Function short form (IIEF-5);
4. Pelvic Floor Distress Inventory Short Form (PFDI-20);
5. Pelvic Floor Impact Questionnaire Short Form (PFIQ-7);
6. Pelvic Organ Prolapse/Urinary Incontinence Sexual Questionnaire Short Form (PISQ-12);
7. Fecal Incontinence Quality of Life Scale (FIQL);
8. Fecal Incontinence Severity Index (FISI).

Contactpersonen

Publiek

's-Gravendijkwal 230, room H-187
E. Utomo
Rotterdam 3015 CE
The Netherlands
+31 (0)10 7030241

Wetenschappelijk

's-Gravendijkwal 230, room H-187
E. Utomo
Rotterdam 3015 CE
The Netherlands
+31 (0)10 7030241

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Aged 18 years or older;
2. Urinary incontinence and/or pelvic organ prolapse (stadium 2 or higher) and/or erectile dysfunction and/or fecal incontinence for at least 3 months.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Malignancy;
2. Dementia;
3. Mental retardation.

Onderzoeksopzet

Opzet

Type:	Observationeel onderzoek, zonder invasieve metingen
Onderzoeksmodel:	Parallel
Toewijzing:	N.v.t. / één studie arm
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving gestopt
(Verwachte) startdatum:	01-06-2010
Aantal proefpersonen:	650
Type:	Werkelijke startdatum

Ethische beoordeling

Positief advies

Datum: 28-05-2010

Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL2229
NTR-old	NTR2355
Ander register	METC / CCMO : 2008-376 / NL25341.078.08 ;
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A