

Effect of intravenous recombinant human APC on coagulation and inflammation in house dust mite and lipopolysaccharide induced allergic asthma.

Gepubliceerd: 14-06-2011 Laatste bijgewerkt: 19-03-2025

Intravenously administered rhAPC has protective effects on HDM-LPS induced allergic lung inflammation.

Ethische beoordeling	Positief advies
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON22425

Bron

Nationaal Trial Register

Verkorte titel

Achilles-study

Aandoening

Allergic asthma, house dust mite allergy, lung inflammation, coagulation

Ondersteuning

Primaire sponsor: Center for Experimental and Molecular Medicine, Academic Medical Center - University of Amsterdam

Overige ondersteuning: Dutch Asthma Foundation

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Influx, differentiation and possible phenotype differences of inflammatory cells.

Toelichting onderzoek

Achtergrond van het onderzoek

Allergic lung inflammation is associated with reduced bronchoalveolar levels of endogenous activated protein C (APC). The biological effects of APC are pleiotropic, and can be roughly divided in anticoagulant and cytoprotective effects. Recombinant human Activated Protein C (rhAPC) has been shown to decrease inflammation and is known for its capability to decrease mortality of patients with severe sepsis. Recent evidence derived from animal studies, in part from our laboratory, indicates that APC is also beneficial in allergic inflammatory conditions. In this study, we will examine whether intravenous administration of rhAPC is capable to inhibit local inflammation, within a lung subsegment, induced by combined administration of house dust mite (HDM) and lipopolysaccharide (LPS) in asthma patients. LPS is a relevant in this context because it is abundant in the natural human environment (and a natural contaminant of HDM) and known to enhance HDM induced allergic inflammation in asthma patients. The primary objective of this study is to determine the effect of intravenously administered rhAPC on HDM-LPS induced allergic lung inflammation.

Doel van het onderzoek

Intravenously administered rhAPC has protective effects on HDM-LPS induced allergic lung inflammation.

Onderzoeksopzet

28 asthma patients will start on intravenous treatment with rhAPC or placebo 4 hours before ($t = -4$ hours) bronchial instillation of HDM/LPS in a lung subsegment and bronchial instillation of saline in a contralateral lung subsegment ($t = 0$ hours). Intravenous treatment with rhAPC or placebo will be continued until 1 hour before initiation of the second bronchoscopy.

Onderzoeksproduct en/of interventie

Intravenous rhAPC treatment and bronchoscopy allergen challenge and lavage. At 05:00 asthma patients will start on intravenous rhAPC treatment or placebo treatment. At 09:00 patients are challenged with housedustmite+LPS in one lungsegment and Saline in a lung segment of the contralateral lung. At 16.00 intravenous treatment is halted and at 17:00 a second bronchoscopy is done to lavage the segments.

Contactpersonen

Publiek

J.D. Boer, de
Academisch Medisch Centrum
Meibergdreef 9
Room F0-117
Amsterdam 1105 AZ
The Netherlands
+31 (0)20 5665247

Wetenschappelijk

J.D. Boer, de
Academisch Medisch Centrum
Meibergdreef 9
Room F0-117
Amsterdam 1105 AZ
The Netherlands
+31 (0)20 5665247

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Intermittent to mild asthmatics between 18 and 45 years of age according to the Global Initiative for Asthma (GINA) criteria;
2. Allergy for HDM documented by a positive RAST and a positive skin prick test;
3. No clinically significant findings during physical examination and hematological and biochemical screening;
4. At spirometry FEV1 more than 70% of predicted value;
5. Able to communicate well with the investigator and to comply with the requirements of the study;
6. Stable asthma without the use of asthma medication 2 weeks prior to the study day;

7. Written informed consent;
8. No current smoking for at least 1 year and less than 10 pack years of smoking history;
9. Both male and female subjects are eligible for the study. Female subjects of child bearing potential will use adequate anti-conceptive precautions and will be tested for pregnancy.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Relevant comorbidity, pregnancy and/or recent surgical procedures;
2. A history of smoking within the last 12 months, or regular consumption of greater than three units of alcohol per day;
3. Exacerbation and/or the use of asthma medication within 2 weeks before start;
4. Administration of any investigational drug within 30 days of study initiation;
5. Donation of blood within 60 days, or loss of greater than 400 ml of blood within 12 weeks of study initiation;
6. History of enhanced bleeding tendency or abnormal clotting test results;
7. History of heparin-induced thrombocytopenia;
8. History of serious drug-related reactions, including hypersensitivity;
9. Inability to maintain stable without the use of asthma medication 2 weeks before start.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blindering:	Dubbelblind
Controle:	Placebo

Deelname

Nederland
Status: Werving gestopt
(Verwachte) startdatum: 01-07-2011
Aantal proefpersonen: 28
Type: Werkelijke startdatum

Ethische beoordeling

Positief advies
Datum: 14-06-2011
Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 35983
Bron: ToetsingOnline
Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL2802
NTR-old	NTR2943
CCMO	NL36336.018.11
ISRCTN	ISRCTN wordt niet meer aangevraagd.
OMON	NL-OMON35983

Resultaten

Samenvatting resultaten

N/A