

monochromatic blue light vs standard light treatment in seasonal complaints

Gepubliceerd: 20-12-2013 Laatste bijgewerkt: 15-05-2024

Seasonal affective disorder, wintertype, according to DSM-IV is depression with a seasonal pattern in which the complaints exist in fall/winter and remission takes place in spring/summer at an almost yearly basis. Epidemiological research in the...

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON22649

Bron

Nationaal Trial Register

Aandoening

Seasonal affective disorder, wintertype, according to DSM-IV is depression with a seasonal pattern in which the complaints exist in fall/winter and remission takes place in spring/summer at an almost yearly basis. Epidemiological research in the Netherlands shows that 3% of the adults suffer from SAD and 8% from winter blues..

Ondersteuning

Primaire sponsor: Philips Consumer Lifestyle/ S. Hermans
P.O. Box 20100
9200 CA Drachten
NL

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

1 - monochromatic blue light vs standard light treatment in seasonal complaints 25-05-2025

Toelichting onderzoek

Achtergrond van het onderzoek

Background of the study:

Seasonal affective disorder, wintertype, according to DSM-IV is depression with a seasonal pattern in which the complaints exist in fall/winter and remission takes place in spring/summer at an almost yearly basis. Epidemiological research in the Netherlands shows that 3% of the adults suffer from SAD and 8% from winter blues..

It has been shown that light treatment is effective, but the etiology of SAD and the working mechanism of light treatment are still unknown. One of the hypotheses is the phase shift hypothesis, which postulated that some biological processes are shifted compared to the 24 h rhythm of the environment. Exposure to bright light can cause a phase shift. If the biological clock is running in phase, SAD complaint can improve..

Recently novel photoreceptors in the eye are discovered. They have no influence on the visual system, but are sensitive for light, especially for light with a short wavelength (blue light). If blue light with a low intensity can have the same effects compared to standard light therapy with a high intensity, then it is possible to simplify the treatment and to incorporate it in the life style of the

Objective of the study:

To investigate the effects of exposure to low intensity monochromatic blue light compared to the effects of standard light therapy in the treatment of SAD and winter blues

Study design:

A treatment study in which in the experimental condition the effects of exposure of low intensity blue monochromatic light is compared to the effects of exposure to standard light treatment in the treatment of SAD and winter blues

Study population:

Patients suffering from SAD are recruited from the SAD outpatient clinic of the UMCG.
Participant suffering from winterblues are recruited by means of advertisements in local newspapers

Intervention (if applicable):

Experimental treatment is exposure to low intensity blue monochromatic light, compared to standard light treatment

Primary study parameters/outcome of the study:

Scores on the SIGH-SAD interviews

Doel van het onderzoek

Seasonal affective disorder, wintertype, according to DSM-IV is depression with a seasonal pattern in which the complaints exist in fall/winter and remission takes place in spring/summer at an almost yearly basis. Epidemiological research in the Netherlands shows that 3% of the adults suffer from SAD and 8% from winterblues..

It has been shown that light treatment is effective, but the etiology of SAD and the working mechanism of light treatment are still unknown. One of the hypotheses is the phase shift hypothesis, which postulated that some biological processes are shifted compared to the 24 h rhythm of the environment. Exposure to bright light can cause a phase shift. If the biological clock is running in phase, SAD complaint can improve..

Recently novel photoreceptors in the eye are discovered. They have no influence on the visual system, but are sensitive for light, especially for light with a short wavelength (blue light). If blue light with a low intensity can have the same effects compared to standard light therapy with a high intensity, then it is possible to simplify the treatment and to incorporate it in the life style of the

Onderzoeksopzet

In a 15 days study design, assessments by means of a standardized structured interviews

(SIG-SAD) at selection/inclusion and at day 1, 8 and 15 (primary outcome)

Daily self ratings (KSS; GSQS; AMS; AD-ACL)

Self ratings at inclusion (expectation, SPAQ and MEQ) and at day 15 (evaluation)

Onderzoeksproduct en/of interventie

1. Patients with Seasonal Affective Disorder, age 18yr. or older

A comparison of the effects of exposure to monochromatic light (blue light) vs standard light treatment

5 consecutive days from 8.00-8.30 a.m. in the clinic, at day 4, 5, 6, 7, 8 of the 15 days study design

2. Subjects with sub-syndromal Seasonal Affective Disorder (winter blues), age 18 yr. or older

A comparison of the effects of exposure to monochromatic light (blue light) vs standard light treatment

5 consecutive days from 8.00-8.20 a.m. or earlier after awakening at home, at day 4, 5, 6, 7, 8 of the 15 days study design

Contactpersonen

Publiek

University Medical Center Groningen, University Center for Psychiatry, PO Box 30 001
W.B. Duijzer
Groningen 9700 RB
The Netherlands
31503613150

Wetenschappelijk

University Medical Center Groningen, University Center for Psychiatry, PO Box 30 001
W.B. Duijzer
Groningen 9700 RB
The Netherlands
31503613150

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

- age between 18-65 yr
- no other treatments for seasonal complaints during the same time
- no traveling to southern counties during study period
- no use of tanning fixtures during study period
- informed consent

1. seasonal affective disorder (SAD), winter type, according to DSM-IV
score of at least 18 on the first 24 items of the SIGH-SAD

2. sub-syndromal seasonal affective disorder (sub-SAD, winterblues) according to the Kasper et al. (1988) criteria:

- SPAQ-GSS score of 8,9 or 10 and at least light seasonal complaints or
- SPAQ-GSS score of 11
score of 12-17 on the first 24 items of the SIGH-SAD

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

other Axis -I disorders according to DSM-IV
acute suicidal risk
use of psychopharmaca or photosensitizing drugs
eye diseases or complaints except aging
diabetes
epilepsy
night shifts

Onderzoeksoepzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Enkelblind
Controle:	Actieve controle groep

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-11-2010
Aantal proefpersonen:	100
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	20-12-2013
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 34402
Bron: ToetsingOnline
Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL4193

Register

NTR-old

CCMO

ISRCTN

OMON

ID

NTR4342

NL33067.042.10

ISRCTN wordt niet meer aangevraagd.

NL-OMON34402

Resultaten

Samenvatting resultaten

N/A