

Efficacy and safety of a single dose of 14.8 MBq/kg (0.4 mCi/kg) ⁹⁰Y-ibritumomab tiuxetan ("Zevalin") in elderly patients with diffuse large B-cell lymphoma and FDG-PET positive partial remission following first-line R-CHOP therapy. A Phase II clinical trial (HOVON 77).

Gepubliceerd: 29-10-2009 Laatste bijgewerkt: 18-08-2022

The hypothesis to be tested is that the efficacy and toxicity of treatment with ⁹⁰Y-ibritumomab tiuxetan meets the expectations as described in the protocol.

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON22777

Bron

Nationaal Trial Register

Verkorte titel

HOVON 77 NHL

Aandoening

Non-Hodgkin's lymphoma (NHL), B-Cell lymphoma

Ondersteuning

Primaire sponsor: Stichting Hemato-Oncologie voor Volwassenen Nederland (HOVON)

P/a HOVON Data Center
Erasmus MC - Daniel den Hoed
Postbus 5201
3008 AE Rotterdam
Tel: 010 7041560
Fax: 010 7041028
e-mail: hdc@erasmusmc.nl

Overige ondersteuning: Koningin Wilhelmina fonds (KWF), Stichting Hemato-Oncologie voor Volwassenen Nederland (HOVON), Bayer Schering Pharma

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Response on FDG-PET (i.e. PET-negative residual masses).

Toelichting onderzoek

Achtergrond van het onderzoek

Study phase: Phase II.

Study objective: Evaluation of efficacy and safety of 90Y-ibritumomab tiuxetan.

Patient population: Patients with Diffuse Large B-Cell lymphoma, CD20-positive, age $\geq$ 60 years and good WHO performance status (0,1,2), with PET-positive PR after R-CHOP induction chemotherapy.

Study design: Prospective, multicenter, open label, non-randomized.

Duration of treatment: Infusion of rituximab followed 1 week later by a second rituximab infusion and a single dose of 90Y-ibritumomab tiuxetan.

Doel van het onderzoek

The hypothesis to be tested is that the efficacy and toxicity of treatment with 90Y-ibritumomab tiuxetan meets the expectations as described in the protocol.

Onderzoeksopzet

At entry, 3 months after treatment, 6 months after treatment or at time of disease

progression, in FU every 3 months during first 2 years, every 6 months during the next 2 years and annually thereafter.

Onderzoeksproduct en/of interventie

Infusion of Rituximab followed one week later by a second Rituximab infusion and a single dose of 90Y-ibritumomab tiuxetan.

Contactpersonen

Publiek

Postbus 7057
J.M. Zijlstra
Amsterdam 1007 MB
The Netherlands
+31 (0)20 4442604

Wetenschappelijk

Postbus 7057
J.M. Zijlstra
Amsterdam 1007 MB
The Netherlands
+31 (0)20 4442604

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Age $\geq$ 60 years old;
2. WHO performance status of 0-2;
3. Life expectancy of at least 3 months;
4. Histologically confirmed CD20 positive Diffuse large B-cell lymphoma (DLBCL), according to the WHO classification;

5. First-line induction treatment with R-CHOP or R-CHOP-like chemotherapy (only CHOP in combination with rituximab; CHOP14 and CHOP21 are both allowed);
6. Partial response on CT-scans after first-line treatment, with measurable disease;
7. PET-positive residual mass;
8. Patient is not eligible for high dose chemotherapy followed by autologous stem cell transplantation;
9. Less than 25% bone marrow involvement at the end of first-line treatment during PR analysis (measurement in a representative bone marrow biopsy);
10. Absolute neutrophil count (ANC) $\geq 1.5 \times 10^9/l$;
11. Hemoglobin (Hb) ≥ 6 mmol/l;
12. Platelets $\geq 150 \times 10^9/l$;
13. Written informed consent obtained according to local guidelines.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Hypoplastic bone marrow at biopsy;
2. Prolonged pancytopenia during induction chemotherapy and delayed courses during R-CHOP induction (more than two weeks delay due to insufficient bone marrow reserve);
3. Known hypersensitivity to murine antibodies or proteins;
4. Significant splenomegaly;
5. Patients with abnormal liver function (total bilirubin $> 2.0 \times$ ULN);
6. Patients with abnormal renal function (serum creatinine $> 2.0 \times$ ULN);
7. Presence of CNS involvement by NHL;
8. Presence of any other active neoplasms or history of prior malignancy, except non-melanoma skin tumours or stage 0 (in situ) cervical carcinoma during the past 5 years;
9. More than one prior R-CHOP or R-CHOP-like chemotherapy regimen for DLBCL;
10. Patients who have received prior external beam radiotherapy to $> 25\%$ of active bone

marrow (involved field or regional);

11. Patients who have received G-CSF or GM-CSF therapy within two weeks prior to study enrollment;

12. Concurrent severe and/or uncontrolled medical condition (e.g. uncontrolled diabetes, congestive heart failure, myocardial infarction within 6 months of study, unstable and uncontrolled hypertension, chronic renal disease, or active uncontrolled infection) which could compromise participation in the study;

13. Patients who have received biologic therapy, immunotherapy, R-CHOP(-like) chemotherapy, surgery, or an investigational drugs less than 4 weeks prior to first day of study treatment or who have not recovered from the toxic effects of such therapy;

14. Patients who have received systemic corticosteroids at doses higher than 20 mg/day prednisolone or equivalent less than 2 weeks prior to 90Y-ibritumomab tiuxetan administration;

15. Known diagnosis of HIV infection;

16. Patients unwilling or unable to comply with the protocol.

Onderzoeksopzet

Opzet

Type: Interventie onderzoek

Onderzoeksmodel: Parallel

Toewijzing: N.v.t. / één studie arm

Controle: N.v.t. / onbekend

Deelname

Nederland

Status: Werving gestart

(Verwachte) startdatum: 05-01-2006

Aantal proefpersonen: 40

Type: Verwachte startdatum

Ethische beoordeling

Positief advies

Datum: 29-10-2009

Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL1969
NTR-old	NTR2086
Ander register	EudraCT number : 2005-003796-20
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A