

Rifampicine vermindert de ontstekingsreactie bij patienten met een (matig) ernstige longontsteking.

Gepubliceerd: 14-12-2012 Laatste bijgewerkt: 19-03-2025

In vitro studies have shown that - in the presence of identical bacterial kill - rifampicin causes reduced release of proinflammatory components of Streptococcus pneumoniae cell wall, lipoteichoic acid (LTA), as compared with other antibiotics...

Ethische beoordeling	Positief advies
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON22991

Bron

Nationaal Trial Register

Verkorte titel

PRISTINE

Aandoening

Community Acquired Pneumonia

Rifampicin

Lipoteichoic Acid

Pneumolysin

inflammatory response

clinical outcome

prognosis

pneumonia

Emergency medicine

hospital

Spoedeisende hulp

ziekenhuis

Thuis opgelopen longontsteking

CURB-65 score

rifampicine
lipoteichoïdzuur
pneumolysine
inflammatoire respons
prognose
klinische uitkomst
pneumonie

Ondersteuning

Primaire sponsor: Prof. dr. J.T. van Dissel
LUMC

Overige ondersteuning: This study was supported by the Virgo consortium, funded by the Dutch government project number FES0908, and by the Netherlands Genomics Initiative (NGI) project number 050-060-452.

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The primary objective is to demonstrate a reduction in lipoteichoic acid release in patients with pneumococcal pneumonia treated with rifampicin and standard treatment as compared to those given standard treatment only. Lipoteichoic acid release will be quantified by measuring lipoteichoic acid in serum and urine. Evaluable patients for Intention-to-Treat analysis are those who met the following criteria: (1) enrollment criteria of pneumonia, with 2 or more points in the CURB65 score, for which a patient is admitted to the hospital, (2) Streptococcus pneumoniae is identified as the cause of pneumonia, (3) received at least one dose of study drugs and (4) outcome is measured at 30 days.

Evaluable patients for Per-Protocol analysis are those who meet the following criteria: (1) enrollment criteria of pneumonia, with 2 or more points in the CURB65 score, for which a patient is admitted to the hospital, (2) Streptococcus pneumoniae is identified as the cause of pneumonia, (3) received the study drug for 48 hours and (4) outcome is measured at 30 days.

Toelichting onderzoek

Achtergrond van het onderzoek

N/A

Doel van het onderzoek

In vitro studies have shown that – in the presence of identical bacterial kill – rifampicin causes reduced release of proinflammatory components of *Streptococcus pneumoniae* cell wall, lipoteichoic acid (LTA), as compared with other antibiotics including benzylpenicillin. Because the inflammatory response to these proinflammatory components determines the intensity of the host immune reaction and, by consequence, collateral tissue damage in infection, a reduced inflammatory response in pneumococcal infection, e.g., community acquired pneumonia, may reduce damage to the lungs and severity of disease in pneumonia.

We will try to demonstrate a reduction in lipoteichoic acid release in patients with pneumococcal pneumonia treated with rifampicin and standard treatment as compared to those given standard treatment only.

Onderzoeksopzet

Assessments of clinical response will be performed continuously during the treatment period by the attending physician, 1-3 days after admission by the clinical researchers, 30 days after start of therapy by a visit of the clinical researchers and 90 days post therapy by a telephone call.

Besides normal diagnostic procedures (sputum culture, blood culture, routine laboratory tests), extra samples are taken:

1. At inclusion: 1 throat swabs, 1 rectum swabs, a urine sample and 4 ml of EDTA blood for biomarker assay and 4 ml of Natriumheparin blood for LTA measurement and 2 ml of blood (in PAXgene RNA tubes);
2. At 2, 4, 8, 16, 24 and 48 hours after inclusion: 8 ml of blood and a urine sample (10ml); at 24 hours we'll also collect 2ml of blood in PAXgene tubes;
3. At day 30: 10 ml of blood, a urine sample (10 ml) and a rectum swab.

Onderzoeksproduct en/of interventie

After informed consent, patients will be randomized (2:1) on the emergency department.

The intervention group (2/3) receives rifampicin 600 mg b.i.d. in the first 48 hours + a beta-lactam antibiotic (mostly penicillin) with or without ciprofloxacin.

The control group (1/3) receives a beta-lactam antibiotic (mostly penicillin) with or without ciprofloxacin.

The choice whether or not to add ciprofloxacin is described in the Dutch SWAB guidelines (<http://www.swab.nl/richtlijnen>).

Contactpersonen

Publiek

Afdeling infectieziekten, C5-P
LUMC
Albinusdreef 2
Postbus 9600
G.H. Groeneveld
Leiden 2300 RC
The Netherlands

Wetenschappelijk

Afdeling infectieziekten, C5-P
LUMC
Albinusdreef 2
Postbus 9600
G.H. Groeneveld
Leiden 2300 RC
The Netherlands

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Patient aged 18 years or above;
2. Hospitalization for community acquired pneumonia with CURB65 score \geq 2.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Known allergy to rifampicin or other rifamycins;
2. Haemolytic anaemia or thrombopenia as side effect of rifampicin in medical history;
3. Liver failure;
4. Use of voriconazol or protease inhibitors.

Note: Patients using other drugs that influence cytochrome P450 3A4 are not excluded. This is an open label trial, all doctors are aware when their patient is treated with rifampicin (or not). If necessary, proper adjustments of concomitant medications can be made (e.g. with oral anticoagulants, ciclosporin). A short period of lower levels of drugs that are prescribed for long-term effects (e.g. antihypertensive drugs or glucose lower medication) will not

interfere with long term outcome. In case female patients in reproductive age use oral contraceptives, other contraceptive measures will be advised after discharge from hospital.

Treatment with rifampicin is for maximum of 48 hours. The enzyme inducing effect is only limited.

The College ter Beoordeling van Geneesmiddelen indicates only voriconazol and protease inhibitors as a real contra-indication.

5. Female patients who are pregnant

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	Actieve controle groep

Deelname

Nederland
Status: Werving gestopt
(Verwachte) startdatum: 07-01-2013
Aantal proefpersonen: 40
Type: Werkelijke startdatum

Ethische beoordeling

Positief advies
Datum: 14-12-2012
Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 39089
Bron: ToetsingOnline
Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL3586
NTR-old	NTR3751
CCMO	NL40521.058.12
ISRCTN	ISRCTN wordt niet meer aangevraagd.
OMON	NL-OMON39089

Resultaten

Samenvatting resultaten

N/A