

A prospective, open-label, multicenter randomized phase-II trial to evaluate the efficacy and safety of a sequential regimen of obinutuzumab (Gazyvaro) followed by obinutuzumab and venetoclax, followed by either standard venetoclax maintenance or MRD guided venetoclax maintenance in first-line patients with CLL and unfit for FCR-like regimens

Gepubliceerd: 27-09-2016 Laatste bijgewerkt: 18-08-2022

Ethische beoordeling	Niet van toepassing
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON23205

Bron

NTR

Verkorte titel

HOVON 139 CLL

Aandoening

Chronic Lymphocytic Leukemia (CLL)

Ondersteuning

Primaire sponsor: HOVON data center

Overige ondersteuning: HOVON, Abbvie, Roche

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

To separately study the efficacy, defined as MRD negative bone marrow and no progression according to the IWCLL criteria, of the two arms of the study of either venetoclax maintenance or MRD-guided venetoclax maintenance after sequential regimens of obinutuzumab (pre-induction) followed by 6 cycles obinutuzumab with venetoclax and 6 cycles of venetoclax (induction) in first-line patients with CLL and unfit for FCR-like regimens.

Toelichting onderzoek

Achtergrond van het onderzoek

Background of the study:

With current therapy, progression free survival of CLL in patients unfit for FCR is around 2 years. Venetoclax treatment, especially when initially combined with an anti-CD20 monoclonal antibody (mAb) has high efficacy and in contrast to kinase inhibitors, has the potency to result in MRD-negative disease status, which possibly allows drug discontinuation. Obinutuzumab has the potency to debulk and therefore when used prior to venetoclax might efficiently prevent the occurrence of tumor lysis syndrome (TLS).

Study design:

A prospective, multicenter, open-label, randomized phase -II trial.

Study population:

First-line patients with CLL and unfit for FCR-like regimens.

Intervention

After pre-induction with obinutuzumab, patients will receive induction treatment with obinutuzumab and/or venetoclax followed by 1 year maintenance with venetoclax (arm A) or MRD guided maintenance with venetoclax (Arm B)

Primary study parameters/outcome of the study

MRD negative bone marrow after maximum 24 cycles of (planned) venetoclax and no progression according to IWCLL criteria at any earlier timepoint.

Onderzoeksopzet

- ◆ Before enrollment: within 28 days before registration, as specified in 10.2
- ◆ After each cycle
- ◆ Induction cycle 1: Before start venetoclax
- ◆ Weekly during venetoclax ramp up in induction cycle 1
- ◆ 15 months after randomization
- ◆ During follow up every 3 months until 3 years after registration. Therafter every 6 months until 7 years after registration or until progression, whatever comes first.

Onderzoeksproduct en/of interventie

After induction treatment venetoclax maintenance will be given for either 1 year (arm A) of or until MRD-negativity with a maximum of 1 year (arm B)

Contactpersonen

Publiek

Academic Medical Center, Amsterdam

Secretariaat Hematologie F4-224

Meibergdreef 9
A.P. Kater
Amsterdam 1105 AZ
The Netherlands
+31 (0)20 5665785

Wetenschappelijk

Academic Medical Center, Amsterdam

Secretariaat Hematologie F4-224

Meibergdreef 9
A.P. Kater
Amsterdam 1105 AZ
The Netherlands
+31 (0)20 5665785

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

- Diagnosis of symptomatic CLL (according to IWCLL guidelines, including minimal required markers (CD5/CD19/CD23 triple positive with light chain restriction))

Patients without prior treatment for CLL (Corticoid treatment administered due to necessary immediate intervention is allowed; within the last 10 days before start of study treatment only dose equivalents of maximum 20 mg prednisolone are permitted);

- ◆ Patients aged ≥ 18 years, not fit for FCR-like regimens, according to the treating physician;
- ◆ Able to adhere to the study visit schedule and other protocol requirements;
- ◆ WHO performance status of ≤ 2 (see appendix C);
- ◆ Laboratory test results within these ranges: - absolute neutrophil count $\geq 1.0 \times 10^9/l$ and platelet count $\geq 50 \times 10^9/l$, unless due to bone marrow infiltration, - creatinine clearance ≥ 45 ml/min (using 24-hour creatinine clearance or modified Cockcroft–Gault equation (see appendix E) - total bilirubin $\leq 1.5 \times$ ULN unless considered due to Gilbert's syndrome, - transaminases $\leq 3 \times$ ULN;
- ◆ Negative serum or urine pregnancy test within 28 days prior to registration (all females of childbearing potential);
- ◆ Written informed consent
- ◆ Patient is capable of giving informed consent

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- ◆ Current inclusion in other clinical trials
- ◆ Intolerance of exogenous protein administration;
- ◆ History of severe allergic or anaphylactic reactions to humanized or murine monoclonal antibodies. Known sensitivity or allergy to murine products.
- ◆ Positive hepatitis serology (serology testing required at screening), as follows:
 - Hepatitis B virus (HBV): Patients with positive serology for hepatitis B defined as positivity for hepatitis B surface antigen (HBsAg) or hepatitis B core antibody (anti-HBc).
 - Hepatitis C virus (HCV): Patients with positive hepatitis C serology unless HCV- (RNA) is confirmed negative.
- ◆ HIV positive patients;
- ◆ Active fungal, bacterial, and/or viral infection that requires systemic therapy; Note: active controlled as well as chronic/recurrent infections are at risk of reactivation/infection during treatment with obinutuzumab and/or venetoclax);
- ◆ Vaccination with a live vaccine a minimum of 28 days prior to registration.
- ◆ Use of any other experimental drug or therapy within 28 days of baseline;
- ◆ Concurrent use of other anti-cancer agents or treatments;
- ◆ History of prior malignancy, except for conditions as listed below if patients have recovered from the acute side effects incurred as a result of previous therapy:
 - Malignancies surgically treated with curative intent and with no known active disease present for ≥ 3 years before randomization
 - Adequately treated non-melanoma skin cancer or lentigo maligna without evidence of disease
 - Adequately treated cervical carcinoma in situ without evidence of disease
- ◆ Severe cardiovascular disease (arrhythmias requiring chronic treatment, congestive heart failure or symptomatic ischemic heart disease) (CTCAE grade III-IV, see appendix D);
- ◆ Severe pulmonary dysfunction (CTCAE grade III-IV, see appendix D);

- ◆ Severe neurological or psychiatric disease (CTCAE grade III-IV, see appendix D);
- ◆ Concurrent severe and/or uncontrolled medical condition (e.g. uncontrolled diabetes, hypertension, hyperthyroidism or hypothyroidism etc.)
- ◆ Women who are pregnant or lactating;
- ◆ Fertile men or women of childbearing potential unless: (1). surgically sterile or ≥ 2 years after the onset of menopause (2). willing to use a highly effective contraceptive method (Pearl Index <1) such as oral contraceptives, intrauterine device, sexual abstinence or barrier method of contraception in conjunction with spermicidal jelly during study treatment and in female patients for 18 months after end of antibody treatment and male patients for 6 months after end of treatment.
- ◆ Any psychological, familial, sociological and geographical condition potentially hampering compliance with the study protocol and follow-up schedule.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	Geneesmiddel

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-12-2016
Aantal proefpersonen:	70
Type:	Verwachte startdatum

Ethische beoordeling

Niet van toepassing	
Soort:	Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL5871
NTR-old	NTR6043
Ander register	2015-004985-27 : HO139 CLL

Resultaten

Samenvatting resultaten

N/A