Ambulatory monitoring of postural instability and gait disorder in Parkinson's Disease.

Gepubliceerd: 17-10-2009 Laatst bijgewerkt: 18-08-2022

Presence of postural instability and gait dissorder PIGD (gait disorders, balance impairment, falls, fal-related injuries and FOG) are an important and disabling feature of idiopathic Parkinson's disease (PD) where the level of functional ability is...

Ethische beoordeling	Niet van toepassing
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Observationeel onderzoek, zonder invasieve metingen

Samenvatting

ID

NL-OMON23798

Bron NTR

Verkorte titel

AMPIGD PD

Aandoening

Ambulatory moniotring Parkinson's Disease Kinematics PIGD

Ondersteuning

Primaire sponsor: Dr. J.P.P. van Vugt

j.vanvugt@mst.nl Medisch Spectrum Twente Afdeling neurologie Postbus 50000 7500 KA Enschede +31 (0)53-487 2841

1 - Ambulatory monitoring of postural instability and gait disorder in Parkinson's D \dots 4-05-2025

Dr. Ir. T. Heida Function: Assistant professor Biomedical Signals and Systems University of Twente Drienerlolaan 5 7522 NB Enschede Building: Zuidhorst 226 +31 (0)53 489 2759

S. Sharifi s.sharifi@student.utwente.nl MSc. Student Technical Medicine Medical Signalling +31 (0)53 4873064

Overige ondersteuning: Dr. J.P.P. van Vugt

j.vanvugt@mst.nl Medisch Spectrum Twente Afdeling neurologie Postbus 50000 7500 KA Enschede +31 (0)53-487 2841

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The subjects walk in a defined environment with six sensors and all activities will be filmed in order to correlate postural disability and gait disorders with the signals registrated by the sensors. The characteristics in gait are to be detected that cause PIGD and the corresponding signal. From this data several features are extracted that would be able to detect and classify PIGD.

Furthermore, with the appointed features an optimal and minimal sensor configuration (amount and position of sensors) will be determined in which these features are still effective in detection and classification.

Toelichting onderzoek

Achtergrond van het onderzoek

Subjects are required to perform a predefined walking trail along a total of 75 m leveled

2 - Ambulatory monitoring of postural instability and gait disorder in Parkinson's D \ldots 4-05-2025

walkway at self-selected preferred speeds. The measurements are subdivided in three sections which will be monitored with 6 sensors including a walking trajectory where tasks are performed which can elicit their walking disabilities and short while free walking (3 min). Furthermore the standard physiotherapy examination performed for Parkinson screening are included in the measurement. At last the UPDRS scale (part III) is obtained. Two questionnaires are asked to be filled in, the Falls efficacy scale and freezing of gait guestionnaire. These measurements will be performed in a defined gym with an even floor at the physiotherapy department of hospital. Patients will be tested during their subjective on/off phase, using their regular medication regimen. The subjects will be summoned for a hospital visit to perform the gait trajectory, the measurement will take about one hour. The aim is to extract features out of the data that is registrated that can detect and classify PIGD. In order to assess the performance of the subject reliable in the home environment with the sensor configurations, performances must be studied on controlled and known trajectories in advance. Therefore a trajectory is designed in which the likelihood of observing any axial disabilities is increased. The sensors only register movement and do no harm. If experiencing serious difficulties with the trajectories there is a risk of making missteps, entering a freezing condition or in worst case falling. The subjects are recruited from the Parkinson population at the Medical Spectrum Twente, Enschede, the Netherlands.

Doel van het onderzoek

Presence of postural instability and gait dissorder PIGD (gait disorders, balance impairment, falls, fal-related injuries and FOG) are an important and disabling feature of idiopathic Parkinson's disease (PD) where the level of functional ability is affected and has a significant impact on quality of life. For clinicians, it is currently difficult to adequately estimate the patient's gait and balance problems in their clinical practice. For most Parkinson's patients, the clinically based assessment does not adequately reflect their actual balance during daily life as it is only an impression at a given moment.

Onderzoeksopzet

All measurements will be done in one time.

Quantification of walking is done by accelerometers and gyroscopes. One triaxiale accelerometer (Dynaport), and five triaxiale accelerometer plus triaxiale gyroscope (Xsens) will be fixed on the body in order to obtain the data. The Dynaport will be fixed only on the lower back as the Xsens sensors all are connected to each other and are placed on the lower back, thigh, shank and both feet.

Furthermore the activities are filmed. The assessment of the clinical state is done by clinical tests UPDRS (part III), Timed-Up-and-Go test, 10.

Onderzoeksproduct en/of interventie

N/A

Contactpersonen

Publiek

Postbus 50000 J.P.P. Vugt, van Enschede 7500 KA The Netherlands +31 (0)53-487 2841

Wetenschappelijk

Postbus 50000 J.P.P. Vugt, van Enschede 7500 KA The Netherlands +31 (0)53-487 2841

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Parkinson patients, which are confirmed idiopathic PD diagnosis according to the UK Parkinson's disease society criteria, in stages 1 to 4 in Hoehn and Yahr scale are recruited for the study. The patients must be able to walk independently without walking aids. The agematched controle group will be able to walk without any visible gait problems.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Impairments or diseases other than PD that affect gait or balance (e.g. orthopedic, neurological other than PD);

- 2. Inability to walk multiple short distances, together totaling approximately half an hour;
- 3. Dependent on walking aids;
- 4. Severe dyskinesia;

4 - Ambulatory monitoring of postural instability and gait disorder in Parkinson's D ... 4-05-2025

5. Uncorrected visual disturbance.

Onderzoeksopzet

Opzet

Туре:	Observationeel onderzoek, zonder invasieve metingen
Onderzoeksmodel:	Anders
Toewijzing:	N.v.t. / één studie arm
Blindering:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-12-2009
Aantal proefpersonen:	25
Туре:	Verwachte startdatum

Ethische beoordeling

Niet van toepassing Soort:

Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

5 - Ambulatory monitoring of postural instability and gait disorder in Parkinson's D ... 4-05-2025

In overige registers

Register	ID
NTR-new	NL1938
NTR-old	NTR2056
Ander register	CCMO ABR-formulier : 30293
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A