

Cardiac Care Bridge trial

Gepubliceerd: 06-04-2017 Laatste bijgewerkt: 18-08-2022

With the Cardiac Care Bridge trial, we hypothesize a 12.5% absolute risk reduction on the composite endpoint of first all-cause unplanned hospital readmission or mortality within six months after randomization.

Ethische beoordeling	Positief advies
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON24273

Bron

NTR

Verkorte titel

CCB-trial

Aandoening

Elderly, aged, older, geriatric, functional decline, functional loss, cardiovascular disease, heart failure, acute myocardial infarction, transitional care, disease management, case management, home-based cardiac rehabilitation, readmission, rehospitalization, mortality

Ouderen, geriatrie, functionele achteruitgang, functieverlies, cardiovasculaire aandoeningen, hartfalen, acuut myocardinfarct, transmurale zorg, diseasesmanagement, casemanagement, hartrevalidatie, heropname, overlijden

Ondersteuning

Primaire sponsor: Academic Medical Center (AMC), Amsterdam, The Netherlands

Overige ondersteuning: ZonMw 'From knowledge to Action II program', grant number 520002002

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The incidence of the composite end-point of first all-cause unplanned hospital readmission or mortality.

Toelichting onderzoek

Achtergrond van het onderzoek

After a hospital admission for heart disease, older patients are at high risk of adverse outcomes such as readmission and death. The current treatment in older cardiac patients is focused on disease management while less attention is paid to general healthcare needs. In the Cardiac Care Bridge (CCB) program we aim to examine the effectiveness of a nurse-coordinated transitional intervention including case management, cardiovascular risk management and home-based rehabilitation for older cardiac patients at high risk of functional loss.

Doel van het onderzoek

With the Cardiac Care Bridge trial, we hypothesize a 12.5% absolute risk reduction on the composite endpoint of first all-cause unplanned hospital readmission or mortality within six months after randomization.

Onderzoeksopzet

The primary outcome will be measured at six months. Secondary outcomes will be measured at three months (phone), six months (home visit) and twelve months (phone)

Onderzoeksproduct en/of interventie

Patients admitted to the department of cardiology or cardiac surgery of > 70 years, at high risk of functional loss and admitted > 48 hours are eligible for inclusion. The program combines case management, disease management and home-based cardiac rehabilitation. All participants will receive a comprehensive geriatric assessment (CGA), performed by a cardiac nurse. Participants in the intervention group will receive care based on identified problems from the CGA in three phases.

1. In the clinical phase, an integrated care plan will be established for all participants. The department of geriatrics will be consulted in case of at least 1 identified problem on the psychological domain or minimal 5 identified geriatric problems in general.

2. In the transitional phase, before discharge, a coordinating community care registered nurse (CCRN) visits the participant in the hospital to receive a face-to-face handover from the cardiac nurse.

3. The post-clinical phase consists of four home visits by the CCRN to continue care based on the integrated care plan, including cardiovascular risk management and evaluation of participants' health status. The CCRN works in close collaboration with the physiotherapist, who will perform nine home-based cardiac rehabilitation sessions to improve participants' functional status.

Contactpersonen

Publiek

Postbus 22660
Bianca Buurman
AMC, Kamer F4-108
Amsterdam 1100 DD
The Netherlands
+31 (0)20 5665991

Wetenschappelijk

Postbus 22660
Bianca Buurman
AMC, Kamer F4-108
Amsterdam 1100 DD
The Netherlands
+31 (0)20 5665991

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

- 70 years and older
- Admitted patients to the departments of cardiology or cardiac surgery
- Admission > 48 hours

- High risk of functional decline according to the VMS screening-tool for frailty of the Dutch Safety Management Program (screening on ADL-functioning, fall risk, malnutrition and delirium): score ≥ 2 in patients aged 70-79 years and score ≥ 1 in patients aged 80 years and older
- Mini-Mental State Examination Score (MMSE) ≥ 15
- Per 28 February 2018, patients with an unplanned hospital admission in the previous six months are also eligible for inclusion (independent of the score on the VMS screening tool for frailty)

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- Congenital heart disease
- Terminal illness: defined as a life expectancy of less than three months, for example because of cancer or serious heart failure.
- Transferred from or planned discharge to a nursing home
- Planned discharge to another hospital not participating in this study
- Unable to communicate in Dutch
- Delirium as confirmed by the treating physician

Onderzoeksoepzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blindering:	Enkelblind
Controle:	Geneesmiddel

Deelname

Nederland	
Status:	Werving gestopt

(Verwachte) startdatum: 06-06-2017
Aantal proefpersonen: 500
Type: Werkelijke startdatum

Voornemen beschikbaar stellen Individuele Patiënten Data (IPD)

Wordt de data na het onderzoek gedeeld: Nog niet bepaald

Ethische beoordeling

Positief advies
Datum: 06-04-2017
Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL6169
NTR-old	NTR6316
Ander register	NL55636.018.16 (CCMO) : METC2016_024 (AMC)

Resultaten

Samenvatting resultaten

Verweij L, Jepma P, Buurman BM, et al. The cardiac care bridge program: design of a randomized trial of nurse-coordinated transitional care in older hospitalized cardiac patients

at high risk of readmission and mortality. BMC Health Serv Res 2018;18(1):508-018-3301-9.