

Insulin-induced microvascular activity in patients with essential hypertension: a possible role for angiotensin II AT1-receptor blockers.

Gepubliceerd: 04-03-2008 Laatste bijgewerkt: 18-08-2022

1: Blockade of the angiotensin II AT1-receptor improves the insulin-induced microvascular effects in hypertensive patients. 2: Blockade of the angiotensin II AT1-receptor impairs the insulin-induced microvascular effects in normotensive control...

Ethische beoordeling	Positief advies
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON24297

Bron

Nationaal Trial Register

Verkorte titel

Role of AT1-receptor blockers in insulin-induced vasodilation.

Aandoening

Hypertension

Hypertensie

Ondersteuning

Primaire sponsor: Prof. CDA Stehouwer

Head Department of Internal Medicine

University Hospital Maastricht

P.O. Box 5800

6202 AZ Maastricht

csteh@sint.azm.nl

Overige ondersteuning: CARIM, Cardiovascular Research Institute Maastricht

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

functional recruitment of capillaries in the skin.

Toelichting onderzoek

Achtergrond van het onderzoek

Background: There is a relation between hypertension and insulin resistance, both associated with increased cardiovascular risk. Hypertension and insulin resistance are characterized by dysfunctions in microcirculation, however it is unclear if microcirculation is the link between these two abnormalities. In addition to its actions in mediating glucose uptake, insulin knows several vascular effects. Insulin induces a vasodilatory response by resistance vessels and preterminal arterioles leading to an overall increase in blood flow (glucose) to the muscles. The local activity of the vasoconstrictor angiotensin II is elevated in patients with hypertension. Previous studies show a possible role for angiotensin II in the hypertensive, insulin resistant phenotype, however a mechanism remains unexplained. In this study we hypothesize that blocking the angiotensin II AT1-receptor improves the insulin-induced microvascular dilatation.

Objectives:

1. Does blockade of the angiotensin II AT1-receptor improve the insulin-induced microvascular effects in hypertensive patients.
2. Does blockade of the angiotensin II AT1-receptor impair the insulin-induced microvascular effects in normotensive control subjects?

Study design:

All subjects will bring 3 visits to the AZM. The following interventions will be applied:

- hyperinsulinemic euglycemic clamp (HEC) + placebo
- HEC + irbesartan (600 mg)
- HEC + felodipine ER (20 mg)

Doel van het onderzoek

- 1: Blockade of the angiotensin II AT1-receptor improves the insulin-induced microvascular effects in hypertensive patients.
- 2: Blockade of the angiotensin II AT1-receptor impairs the insulin-induced microvascular effects in normotensive control subjects.

Onderzoeksopzet

Each visit takes approx. 7,5 hr. During all visits 2 catheters will be inserted in the antecubital vein of the lower arms. On one study day (randomly assigned) a set of microcirculation measurements will be performed on $t=-90$ minutes. On all three study days insulin and glucose will be infused on $t=0$ min. After 90 minutes of HEC a set microcirculation measurements will be done, and after these measurements placebo, irbesartan or felodipine will be taken in a single oral dose. 210 minutes after intake ($t=300$ min.) another set of microcirculation measurements will be done. During the study days the heart rate and blood pressure will be monitored and blood samples will be taken. The interventions will be randomly assigned. One week is scheduled between each visit.

Onderzoeksproduct en/of interventie

Hypertensive subjects will be asked to discontinue the intake of antihypertensive medication three weeks before the start of the study. All subjects will be asked to start a low salt diet (100mmol/day) 7 days prior to the first study day and to collect urine during 24hrs prior to the first study day.

Three study days will be performed (each day lasts 7,5hr). The following measurements will be done:

Microcirculation measurements: 1) perfused capillary density and functional capillary recruitment in the nailfold, visualized by a capillary microscope, 2) endothelium-(in)dependent vasodilation of finger skin microcirculation, evaluated with laser Doppler measurements in combination with iontophoresis of acetyl-choline and sodium nitroprusside, and 3) densities and diameter of arterioles, capillaries and venules in the bulbar conjunctiva, measured with conjunctival microscopy.

Placebo, irbesartan (600mg) and felodipine (20mg) will be ingested orally in a single dose. Insulin is infused in a primed continuous manner at a rate of $50\text{mU}\cdot\text{kg}^{-1}\cdot\text{hr}^{-1}$. Euglycemia will be maintained by adjusting the rate of a 20% D-glucose infusion based on plasma glucose measurements performed at 5 min intervals. During the visit several blood samples will be taken, blood pressure and heart rate will be monitored.

Contactpersonen

Publiek

Department of Internal Medicine
University Hospital Maastricht
P.O. Box 5800

A.J.H.M. Houben
Department of Internal Medicine

Maastricht 6202 AZ
The Netherlands

Wetenschappelijk

Department of Internal Medicine
University Hospital Maastricht
P.O. Box 5800

A.J.H.M. Houben
Department of Internal Medicine

Maastricht 6202 AZ
The Netherlands

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Inclusion criteria hypertensive subjects:

1. 18-60 years
2. Caucasian
3. untreated hypertension >140/90mmHg.

Inclusion criteria normotensive subjects:

1. 18-60 years
2. Caucasian
3. Blood pressure <140/90 mmHg.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Obesity (BMI>27kg/m²)
2. Cardiovascular disease (stroke, coronary artery disease, peripheral vascular disease, heart failure)
3. Impaired glucose tolerance or diabetes mellitus according to the criteria of the ADA
4. Smoking
5. Alcohol use >4U/day
6. Use of medication (antihypertensive drugs, lipid lowering drugs, corticosteroids, NSAIDs)

- 7. Pregnancy
- 8. Wearing contact lenses

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Cross-over
Toewijzing:	N.v.t. / één studie arm
Blinding:	Dubbelblind
Controle:	Placebo

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-03-2008
Aantal proefpersonen:	32
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	04-03-2008
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL1158
NTR-old	NTR1202
Ander register	MEC : MEC 07-2-115
ISRCTN	ISRCTN wordt niet meer aangevraagd

Resultaten

Samenvatting resultaten

N/A