

# Onderzoek naar de werkzaamheid van Caphosol tegen mucositis bij kinderen met kanker.

Gepubliceerd: 15-06-2010 Laatst bijgewerkt: 15-05-2024

The null hypothesis will be that Caphosol does not decrease the number of days of mucositis > grade 1, whereas the alternative hypothesis is that it will decrease the number of days of mucositis > grade 1 with 50%.

|                             |                       |
|-----------------------------|-----------------------|
| <b>Ethische beoordeling</b> | Positief advies       |
| <b>Status</b>               | Werving gestopt       |
| <b>Type aandoening</b>      | -                     |
| <b>Onderzoekstype</b>       | Interventie onderzoek |

## Samenvatting

### ID

NL-OMON24436

### Bron

NTR

### Verkorte titel

Caphosol and mucositis

### Aandoening

Hospitalised children with cancer therapy induced mucositis

## Ondersteuning

**Primaire sponsor:** Universitair Centrum van Groningen, Utrecht, VU en Leiden

**Overige ondersteuning:** eerste geldstroom (geld van ministerie van OC&W aan universiteiten) en derde geldstroom door Eusapharma (farmaceutische industrie)

## Onderzoeksproduct en/of interventie

## Uitkomstmaten

### Primaire uitkomstmaten

Days of mucositis > grade I.

## Toelichting onderzoek

### Achtergrond van het onderzoek

Due to more intensive chemotherapy, overall survival in pediatric oncology patients has increased dramatically over the last decades. With this increased survival, it has become even more important to optimize the quality of life during and after treatment. One of the important side effects in pediatric oncology is chemotherapy or radiotherapy induced oral mucositis. This mucositis is accompanied with decreased oral intake, (severe) pain, often with use of analgesics and hospital admission. Thus, it has a severe impact on quality of life. Coincidental with chemotherapy induced mucositis is therapy-related neutropenia. Neutropenic patients with mucositis are at increased risk for systemic infection. In a case controlled study, patients with mucositis and neutropenia had four times the risk of septicaemia than neutropenic patients without mucositis. Until now, no clinically useful drugs are available to prevent or treat chemotherapy induced oral mucositis. Recently, Caphosol, a  $\text{Ca}^{2+}$  /  $\text{PO}_4^{3-}$  mouth rinse, became available to treat or prevent mucositis. Until now, only one clinical study has been performed which concerned adult bone marrow transplant patients receiving chemotherapy and or radiotherapy. In these patients, Caphosol used as prophylaxis reduced the frequency, intensity and duration of oral mucositis. No information is available about the prophylactic or therapeutic use in children treated with chemotherapy or radiotherapy. Nevertheless, Caphosol is licensed as medical device for the treatment and prophylaxis of oral mucositis for all ages. Therefore, we will study the effect of the therapeutic use of Caphosol in cancer therapy induced mucositis in pediatric oncology patients in 4 academic pediatric oncology hospitals in The Netherlands.

### Doel van het onderzoek

The null hypothesis will be that Caphosol does not decrease the number of days of mucositis > grade 1, whereas the alternative hypothesis is that it will decrease the number of days of mucositis > grade 1 with 50%.

### Onderzoeksopzet

Morbidity measurements:

Experienced nurses will take tests 2 times daily in the morning and the afternoon. The following items will be scored:

1. Mucositis, using the NCI CTCAE mucositis score<sup>3</sup>;
2. Pain score: VAS score<sup>4</sup> and faces score<sup>5</sup>;

3. Pain medication: product, dosage;
4. Use of antibiotics and bloodcultures taken;
5. Use of tube and/or parenteral feeding;
6. Tolerability of the product.

## **Onderzoeksproduct en/of interventie**

### Study design:

Since it is not ethical to ask patients consent to enter the study when they present with mucositis in such a degree that it needs hospital admission and medication, another strategy has been chosen. All patients at risk to develop oral mucositis on basis of their chemotherapy treatment, will be asked to participate in the study, i.e. that they will get either Caphosol or standard treatment when they are eligible for the study. When patients present in the hospital with oral mucositis fulfilling the inclusion criteria for treatment with mouth rinse, they will be randomised and get either study- or placebo mouth rinse.

After randomisation, the patient will get standard supportive care, plus study treatment (Caphosol or NaCl 0.9%, i.e. standard mouth rinse). Standard treatment will be according to the institutes protocol and includes adequate pain management, hydration management to guarantee optimal fluid / hydration state, meaning nasogastric tube feeding when possible, or intravenous (re)hydration with Dextrose, NaCl and KCl (plus TPN if necessary). Oral care includes brushing of the teeth 3 - 4 times per day if possible, and standard care according to local policies. The patients will rinse their mouth with study rinse (Caphosol or NaCl 0.9%) 4 times daily, 1 minute with 50% of the solution (15ml), and than 1 minute with the other half (15ml). If this is not possible (for the smaller children or when it is too painful to get this amount of fluid in the mouth), it is allowed to rinse with smaller portions until 30 ml in 2 minutes is reached. After rinsing, the rinse is to be spit out. The nurse on the ward will note the time and amount of mouthrinse. The alternative study treatment , NaCl 0.9%, also (as Caphosol) has a salty taste and is currently the standard treatment for oral mucositis. Rinse should be used 15 minutes before or after meals and before sleeping time. As long as the patient suffers from mucositis the treatment will be given.

## **Contactpersonen**

## **Publiek**

Posbus 30 001  
W.J.E. Tissing  
Groningen 9700 RB  
The Netherlands  
+31 (0)50 3614213

## **Wetenschappelijk**

Posbus 30 001  
W.J.E. Tissing  
Groningen 9700 RB  
The Netherlands  
+31 (0)50 3614213

## **Deelname eisen**

### **Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)**

1. Age 4-18 years;
2. Cancer therapy induced oral mucositis;
3. Hospitalisation.

### **Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)**

1. Impossibility to rinse the mouth;
2. Previous study participation;
3. Previous Caphosol use.

## **Onderzoeksopzet**

## Opzet

|                  |                       |
|------------------|-----------------------|
| Type:            | Interventie onderzoek |
| Onderzoeksmodel: | Parallel              |
| Toewijzing:      | Gerandomiseerd        |
| Blinding:        | Dubbelblind           |
| Controle:        | Geneesmiddel          |

## Deelname

|                         |                       |
|-------------------------|-----------------------|
| Nederland               |                       |
| Status:                 | Werving gestopt       |
| (Verwachte) startdatum: | 01-07-2010            |
| Aantal proefpersonen:   | 30                    |
| Type:                   | Werkelijke startdatum |

## Ethische beoordeling

|                 |                  |
|-----------------|------------------|
| Positief advies |                  |
| Datum:          | 15-06-2010       |
| Soort:          | Eerste indiening |

## Registraties

### Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 35477  
Bron: ToetsingOnline  
Titel:

### Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

## In overige registers

| Register | ID     |
|----------|--------|
| NTR-new  | NL2250 |

**Register**

NTR-old

CCMO

ISRCTN

OMON

**ID**

NTR2377

NL29501.042.09

ISRCTN wordt niet meer aangevraagd.

NL-OMON35477

## Resultaten

**Samenvatting resultaten**

N/A