

Fusie van cardiale CT en PET scan voor bepalen van normale glucose (suiker) opname rondom een Aorta Kunstklep

Gepubliceerd: 14-06-2016 Laatste bijgewerkt: 18-08-2022

Prosthetic Heart Valve (PHV) endocarditis is complicated by peri-annular extension (abscesses/mycotic aneurysms) in up to 50% of patients and has an in-hospital mortality of approximately 30%. However, with current standard imaging tools such as...

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON24441

Bron

NTR

Verkorte titel

PROSPECTA

Aandoening

PET-CT, CTA, prosthetic heart valve, endocarditis

Ondersteuning

Primaire sponsor: University Medical Center Utrecht, Erasmus Medical Center Rotterdam

Overige ondersteuning: None

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

- Qualification Visual Score for Hypermetabolism (QVSH)

- Standardized Uptake Value (SUV) ratio (quantitative measurement)

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale:

Prosthetic Heart Valve (PHV) endocarditis is complicated by peri-annular extension (abscesses/mycotic aneurysms) in up to 50% of patients and has an in-hospital mortality of approximately 30%. However, with current standard imaging tools such as Echocardiography and Computed Tomography Angiography (CTA), it remains difficult to detect peri-annular extension, which is an indication for urgent surgery in order to reduce mortality. Furthermore, if peri-annular abnormalities are found with imaging, differentiation between active and non-active inflammatory tissue is impossible, because of the lack of metabolic information. Hybrid imaging with combined anatomical information by CTA and metabolic information by FDG-PET may provide this additional information. However, normal baseline FDG uptake at different time points after aortic PHV implantation is unknown and obligatory for correct interpretation of CTA/PET scans in PHV patients.

Objective: To determine normal FDG-uptake around PHV's in aortic position.

Study design: Prospective multi-centre cross-sectional study

Methods: 18-Fluorine FDG-PET is performed to assess uptake around the PHV after fusion with the CTA. FDG uptake is scored by: 1. Qualification Visual Score for Hypermetabolism (QVSH) which can be: none, mild, moderate or severe. 2. Standardized Uptake Value (SUV) ratio's, defined as the maximum SUV value around the PHV divided by the mean SUV value in the mediastinum.

Study population: The total group consists of patients (≥ 50 years) after uncomplicated PHV implantation in aortic position ($n=75$). The FDG-PET/CTA imaging is performed in the early and late and chronic postoperative episode (group 1, 2 and 3 respectively):

- Early postoperative (group 1, $n=25$): $5 (\pm 1)$ weeks after PHV implantation .
- Late postoperative (group 2, $n=25$): $12 (\pm 2)$ weeks after PHV implantation .
- Chronic postoperative (group 3, $n=25$): $12 (\pm 2)$ months after PHV implantation.

We will only include patients in group 3 in case FDG uptake around the PHV in group 2 results in moderate or severe FDG uptake according to the QVSH score in 1 or more patients.
Of note: patients cannot be included in two groups.

Main study parameters: 18F-FDG/PET baseline uptake measured by the QVSH value and the SUV ratio's around the PHV in early, late and possibly also in the chronic postoperative phase: 5 ($\pm$ 1) weeks, 12 ($\pm$ 2) weeks and 12 ($\pm$ 2) months respectively.

Doel van het onderzoek

Prosthetic Heart Valve (PHV) endocarditis is complicated by peri-annular extension (abscesses/mycotic aneurysms) in up to 50% of patients and has an in-hospital mortality of approximately 30%. However, with current standard imaging tools such as Echocardiography and Computed Tomography Angiography (CTA), it remains difficult to detect peri-annular extension, which is an indication for urgent surgery in order to reduce mortality. Furthermore, if peri-annular abnormalities are found with imaging, differentiation between active and non-active inflammatory tissue is impossible, because of the lack of metabolic information. Hybrid imaging with combined anatomical information by CTA and metabolic information by FDG-PET may provide this additional information. However, normal baseline FDG uptake at different time points after aortic PHV implantation is unknown and obligatory for correct interpretation of CTA/PET scans in PHV patients.

Onderzoeksopzet

cross-sectional

Onderzoeksproduct en/of interventie

PET-CT and CTA

Contactpersonen

Publiek

University Medical Center Utrecht, Department of Radiology
P.O. Box 85500, E01.132

A.M. Harder, den
Utrecht 3508 GA
The Netherlands

Wetenschappelijk

University Medical Center Utrecht, Department of Radiology
P.O. Box 85500, E01.132

A.M. Harder, den
Utrecht 3508 GA
The Netherlands

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Age $\geq$ 50 years

Patients after uncomplicated PHV implantation in aortic position (mechanical and biological PHVs).

Normal routine follow up TTE (standardly performed 5 days after operation). With no signs of obstruction, endocarditis or significant paravalvular leakages.

Group 1: 5 ($\pm$ 1) weeks after PHV implantation (n=25).

Group 2: 12 ($\pm$ 2) weeks after PHV implantation (n=25).

Group 3 : 12 ($\pm$ 2) months after PHV implantation (n=25). Inclusion in group 3: only in case of $\geq$ 1 patients with post-operative moderate/severe uptake (QVSH score) in group 2.

Weight < 110 kg

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Known contrast allergy

Known renal impairment (GFR<60)

Diabetes Mellitus

Mild contractile dysfunction of the left and/or right ventricular (Eyeballing, Ejection fraction <45 %, TAPSE <14 mm)

Active cardiac decompensation

Uncontrolled cardiac arrhythmias

Suspicion on active endocarditis

Previous participation in scientific studies using radiation.

(Possible) pregnancy in pre-menopausal women above 50 years not on reliable birth control therapy. Other contraindications for contrast use according the standard daily clinical routine according to the protocol "Protocol preventie contrastreactie en contrastnefropathie" by the department of radiology UMCU or EMC

Use of pericardial patches and re-operation of aortic PHV in past medical history

Contraindication for Computed Tomography Angiography according the standard daily clinical routine

Refusal to be informed about potential additional CTA or FDG-PET findings

If already included in group 1 (early post operative phase), patients cannot be included in group 2 or 3 (late post operative phase).

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Niet-gerandomiseerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	21-05-2013
Aantal proefpersonen:	75
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	14-06-2016
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL5478
NTR-old	NTR5895
Ander register	NL42743.041.12 : 12-633

Resultaten