

Pegvisomant-toevoeging bij patiënten met acromegalie: Effect op symptomen, kwaliteit van leven en insulinegevoeligheid.

Gepubliceerd: 22-08-2011 Laatst bijgewerkt: 18-08-2022

1. Addition of weekly pegvisomant administrations improves quality of life; 2. Addition of weekly pegvisomant administrations improves insulin sensitivity.

Ethische beoordeling	Niet van toepassing
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON24812

Bron

Nationaal Trial Register

Aandoening

Acromegaly, pituitary disease, normal IGF1, somatostatin analogues, Pegvisomant

Ondersteuning

Primaire sponsor: Erasmus University MC

Dept. of Medicine

PO Box 2040

3000 CA Rotterdam NL

Tel: +31 10 703 2862

Overige ondersteuning: Placebo and Pegvisomant will be supplied by Pfizer. Additional costs will paid by the department of internal medicine.

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Change in the Quality of Life over 16 weeks as assessed by AcroQoL and PASQ.

Toelichting onderzoek

Achtergrond van het onderzoek

Randomized double blind multi-centre study of the effects on low-dose pegvisomant treatment in acromegalic subjects in whom the IGF-I levels has been normalized by long-acting somatostatin analogs.

Primary Study Objective:

To assess the efficacy and safety of the co-administration of low-dose pegvisomant (40 mg, administered via subcutaneous injection given once a week) and long-acting somatostatin analogs (administered once monthly) on the Quality of Life over 16 weeks in 60 acromegalic patients.

The primary endpoint will be the change in the AcroQoL-physical score at the end of the treatment period.

Secondary Study Objectives:

To assess the effect of low-dose pegvisomant co-administration on: Quality of life and insuline sensitivity.

1. AcroQoL and PASQ;
2. Oral glucose loading;
3. Total body water / body weight;
4. Blood pressure;
5. HbA1c;
6. BNP levels;

7. Ring-size;
8. IGF-I levels;
9. Safety based on:
 - A. Adverse events, clinical examination, vital signs;
 - B. Glucose tolerance;
 - C. Standard hematology and biochemistry, including liver function tests.

Doel van het onderzoek

1. Addition of weekly pegvisomant administrations improves quality of life;
2. Addition of weekly pegvisomant administrations improves insulin sensitivity.

Onderzoeksopzet

Efficacy will be assessed as change between prestudy visit and visit at 16 weeks.

Onderzoeksproduct en/of interventie

Weekly sc administration of Pegvisomant (40 mg) or placebo during 16 weeks.

Contactpersonen

Publiek

Erasmus University MC
Dept. of Medicine
PO Box 2040
S.J.C.M.M. Neggers
Rotterdam 3000 CA
The Netherlands
+31 (0)10 7032862

Wetenschappelijk

Erasmus University MC
Dept. of Medicine
PO Box 2040

S.J.C.M.M. Neggers
Rotterdam 3000 CA
The Netherlands
+31 (0)10 7032862

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Provision of written informed consent prior to any study related procedures;
2. Male or female aged between 18 and 75 years inclusive;
3. The patient must have had documentation supporting the diagnosis of acromegaly based on elevated GH and/or IGF-1 levels;
4. The patient is treated with lanreotide Autogel or octreotide LAR for at least 6 months and has a serum IGF-1 level above the 60th percentile and below ULN, 28 days after the last injection.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Has undergone pituitary surgery or radiotherapy within 6 months prior to study entry;
2. It is anticipated that the patient will receive pituitary surgery or radiotherapy during the study;
3. Has a history of hypersensitivity to lanreotide, octreotide or pegvisomant or drugs with a similar chemical structure;
4. Has already been treated with a somatostatin analogue associated with pegvisomant;
5. Has received a dopamine agonist within 6 weeks prior to study entry;
6. Has been treated with any unlicensed drug within the last 30 days before study entry;
7. Has abnormal hepatic function at study entry (defined as AST, ALT, gGT, alkaline phosphatase, or total bilirubin above 2 ULN);
8. Is at risk of pregnancy or is lactating. Females of childbearing potential must provide a

negative pregnancy test within 5 days before the start of the study and must be using contraception. Non-childbearing potential is defined as post-menopause for at least one year, surgical sterilization or hysterectomy at least three months before the start of the study;

9. Has a history of, or known current, problems with alcohol or drug abuse;
10. Has a mental condition rendering the subject unable to understand the nature, scope and possible consequences of the study, and/or evidence of an uncooperative attitude;
11. Has abnormal baseline findings, any other medical condition(s) or laboratory findings that, in the opinion of the investigator, might jeopardize the subject's safety or decrease the chance of obtaining satisfactory data needed to achieve the objective(s) of the study;
12. Renal insufficiency, clearance < 60 ml/min;
13. Participation in a clinical trial in the last 6 months.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Dubbelblind
Controle:	Placebo

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-12-2011
Aantal proefpersonen:	60
Type:	Verwachte startdatum

Ethische beoordeling

Niet van toepassing	
Soort:	Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL2886
NTR-old	NTR3032
Ander register	:
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

Neggers SJ, van Aken MO, de Herder WW, Feelders RA, Janssen JA, Badia X, Webb SM, van der Lely AJ (2008) Quality of life in acromegalic patients during long-term somatostatin analog treatment with and without pegvisomant. The Journal of clinical endocrinology and metabolism 93 (10):3853-3859.

Biermasz NR, Pereira AM, Smit JW, Romijn JA, Roelfsema F (2005) Morbidity after long-term remission for acromegaly: persisting joint-related complaints cause reduced quality of life. The Journal of clinical endocrinology and metabolism 90 (5):2731-2739.

Bonapart IE, van Domburg R, ten Have SM, de Herder WW, Erdman RA, Janssen JA, van der Lely AJ (2005) The 'bio-assay' quality of life might be a better marker of disease activity in acromegalic patients than serum total IGF-I concentrations. European journal of endocrinology / European Federation of Endocrine Societies 152 (2):217-224.