

# SO-HIP study

Gepubliceerd: 01-04-2016 Laatste bijgewerkt: 19-03-2025

After hip fracture, 20-90% of older individuals experience new disabilities in activities of daily living (ADL) that affect their ability to remain living independently at home. Fear of falling hinder older persons to perform ADLs needed for good...

<b>Ethische beoordeling</b>	Positief advies
<b>Status</b>	Werving gestart
<b>Type aandoening</b>	-
<b>Onderzoekstype</b>	Interventie onderzoek

## Samenvatting

### ID

NL-OMON24960

### Bron

Nationaal Trial Register

### Aandoening

hip fracture, fear of falling

### Ondersteuning

**Primaire sponsor:** Academic Medical Center (AMC), University of Amsterdam (UVA), The Netherlands

**Overige ondersteuning:** Fonds Nuts OHRA  
NWO

### Onderzoeksproduct en/of interventie

### Uitkomstmaten

#### Primaire uitkomstmaten

patients' perceived daily functioning, assessed with the performance rating of the Canadian Occupational Performance Measure (COPM)

# Toelichting onderzoek

## Achtergrond van het onderzoek

Six nursing homes will be randomized in a three-arm stepped wedge cluster randomized trial. All nursing homes will initially provide standard care. At designated time points, nursing homes, successively and in random order, will cross over to the provision of OT and at the next time point, to sensor monitoring-enhanced OT. A total of 360 older individuals, previously living alone in the community, who after a hip fracture were admitted to a geriatric rehabilitation ward for a short-term rehabilitation, will be enrolled.

Individuals in the first intervention group (OTc) will participate in an OT rehabilitation program based on cognitive behavioral therapy (CBT) principles. In the sensor monitoring group, sensor monitoring is added to the OT CBT (OTcsm).

Participants will receive a sensor monitoring system consisting of (i) an activity monitor during nursing home stay, (ii) a sensor monitoring system at home and a (iii) a web-based feedback application. These components will be embedded in the OT. The OT consists of a weekly session with an occupational therapist during the nursing home stay followed by four home visits and four telephone consultations. The primary outcome is patient-perceived daily functioning at 6 months, assessed using the Canadian Occupational Performance Measure (COPM).

## Doel van het onderzoek

After hip fracture, 20-90% of older individuals experience new disabilities in activities of daily living (ADL) that affect their ability to remain living independently at home. Fear of falling hinder older persons to perform ADLs needed for good recovery. Increasing self-efficacy beliefs can reduce fear of falling and can help increase physical activity needed to recover. By goal setting, better insight in the ADLs of older persons performance an recovery after hip fracture might be enhanced. Sensor monitoring can be used as a coaching tool. Currently it is unclear how effective sensor monitoring embedded in an occupational therapist (OT)rehabilitation program is in the recovery of ADLs compared by coaching by an OT alone or care as usual.

We expect that the use of sensor monitoring, embedded in an OT-rehabilitation program in nursing homes and a follow-up at the home of the participant leads to more improvement in ADLs of older persons after a hip fracture compared to the OT coaching without sensor monitoring and usual care. This results in enhanced participation in daily activities, improvement of sense of safety and improved self-rated health related quality of life.

## Onderzoeksopzet

four moments of measurements for all patients:

- first week nursing home (T0)

- before discharge from the nursing home to home (T1)
- T3, post- intervention, 4 months after the start of the geriatric rehabilitation
- 6 months after the start of the geriatric rehabilitation (T6)

### **Onderzoeksproduct en/of interventie**

1. care as usual (multidisciplinary rehabilitation)
2. care as usual and coaching by an OT using cognitive behavioral treatment principles concerning falls
3. care as usual and sensor monitoring added to the coaching by an OT

## **Contactpersonen**

### **Publiek**

Amsterdam University of Applied Science - Research Group Occupational Therapy , ACHIEVE,  
Centre of Applied Research, Faculty of Health,  
Margriet Pol  
Tafelbergweg 51 Po Box 2557  
Amsterdam 1000 CN  
The Netherlands

### **Wetenschappelijk**

Amsterdam University of Applied Science - Research Group Occupational Therapy , ACHIEVE,  
Centre of Applied Research, Faculty of Health,  
Margriet Pol  
Tafelbergweg 51 Po Box 2557  
Amsterdam 1000 CN  
The Netherlands

## **Deelname eisen**

### **Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)**

Patients are:

- admitted to a geriatric rehabilitation ward in a nursing home after hip surgery
- 65 years and older
- living alone in the community or in a senior residence
- have a minimal-mental state examination of 15 or higher
- have an indication for a short term rehabilitation

### **Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)**

- are terminally ill
- are waiting for a permanent place in a nursing home
- do not give informed consent

## **Onderzoeksopzet**

### **Opzet**

Type:	Interventie onderzoek
Onderzoeksmodel:	Cross-over
Toewijzing:	Gerandomiseerd
Blinding:	Enkelblind
Controle:	Geneesmiddel

### **Deelname**

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-04-2016
Aantal proefpersonen:	280
Type:	Verwachte startdatum

## Ethische beoordeling

Positief advies

Datum: 01-04-2016

Soort: Eerste indiening

## Registraties

### Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 43760

Bron: ToetsingOnline

Titel:

### Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

### In overige registers

Register	ID
NTR-new	NL5610
NTR-old	NTR5716
CCMO	NL54132.018.15
OMON	NL-OMON43760

## Resultaten