

# **Arginine suppletie om de reperfusie schade te verminderen en doorbloeding te verbeteren na vrije weefsel transplantaties.**

Gepubliceerd: 13-01-2009 Laatst bijgewerkt: 18-08-2022

L-arginine, a precursor of NO, reduces ischemia-reperfusion injury and increases microcirculatory bloodflow. This will lead to a reduction of complications in free flap surgery.

<b>Ethische beoordeling</b>	Positief advies
<b>Status</b>	Werving gestopt
<b>Type aandoening</b>	-
<b>Onderzoekstype</b>	Interventie onderzoek

## **Samenvatting**

### **ID**

NL-OMON25004

### **Bron**

Nationaal Trial Register

### **Verkorte titel**

N/A

### **Aandoening**

reperfusion injury, Arginine, free flap surgery

### **Ondersteuning**

**Overige ondersteuning:** Plastic Surgery Department, Maastricht University Center of Health. Dutch Burns Foundation

### **Onderzoeksproduct en/of interventie**

### **Uitkomstmaten**

#### **Primaire uitkomstmaten**

Partial flap loss (result of insufficient bloodflow).

## Toelichting onderzoek

### Achtergrond van het onderzoek

Partial flap loss due to microvascular failure is first initiated by the incapability of the vascular pedicle to provide sufficient microvascular perfusion in distal segments of the flap. In addition, in free flap surgery the entire flap is affected by ischemia reperfusion injury. The distal segments in free flap surgery are particular vulnerable for ischemia reperfusion injury, and are thus primarily affected by ischemia reperfusion injury which may lead to distal partial flap loss.

Nitric oxide (NO) has been the focus in an extensively amount of studies regarding its use in reducing the IR-injury. It is widely accepted that the L-Arginine-Nitric Oxide pathway plays a pivotal role in the pathophysiology of IR-injury. L-Arginine, the sole precursor of NO, can be metabolized in NO and citrulline by a family of tree isoforms of NO synthase (NOS).

Endothelial NOS (eNOS) and neuronal NOS (nNOS), these are mainly constitutively expressed by respectively endothelial cells and neuronal cells. Expression of inducible NOS (iNOS) is induced by inflammatory mediators and is mainly expressed in leucocytes. L-Arginine has shown to scavenge free radicals which are expressed during reperfusion. Its end product NO is an important and potent vasodilatator and prevents aggregation and activation of neutrophils and platelets. Furthermore NO concentrations are reduced during ischemia and production remains low after reperfusion. Therefore increasing NO production by stimulating the L-Arginine-NO pathway may lessen the severity of IR-injury. In experimental studies intravenous L-Arginine substantially reduces ischemia-reperfusion injury in cutaneous and musculocutaneous flaps. The purpose of this translational study was to establish the possible protective effect of L-arginine on microvascular perfusion and clinical outcome in free flap surgery.

### Doel van het onderzoek

L-arginine, a precursor of NO, reduces ischemia-reperfusion injury and increases microcirculatory bloodflow. This will lead to a reduction of complications in free flap surgery.

### Onderzoeksopzet

1. After flap dissection;
2. ischemia;
3. reperfusion;
4. 1,2,3,4,5 hours after reperfusion;

5. 1 week outpatient clinic;

6. 6 week outpatient clinic.

### **Onderzoeksproduct en/of interventie**

Patients received intravenously one liter of either L-arginine (verum group, 30 g L-arginine-HCl / 1L 0.9% NaCl) or an Alanine (placebo group, 25.2 g Alanine / 1L 0.9% NaCl) during a 24 hour period.

## **Contactpersonen**

### **Publiek**

P. Debeyelaan 25  
R.R.W.J. Hulst, van der  
Maastricht 6202 AZ  
The Netherlands  
+ 31 433875624

### **Wetenschappelijk**

P. Debeyelaan 25  
R.R.W.J. Hulst, van der  
Maastricht 6202 AZ  
The Netherlands  
+ 31 433875624

## **Deelname eisen**

### **Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)**

All patients with free TRAM flap breast reconstruction.

### **Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)**

Previous midline laparotomy.

# Onderzoeksopzet

## Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Dubbelblind
Controle:	Placebo

## Deelname

Nederland	
Status:	Werving gestopt
(Verwachte) startdatum:	01-09-2003
Aantal proefpersonen:	40
Type:	Werkelijke startdatum

## Ethische beoordeling

Positief advies	
Datum:	13-01-2009
Soort:	Eerste indiening

## Registraties

### Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

### Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

## In overige registers

Register	ID
NTR-new	NL688
NTR-old	NTR1626
Ander register	:
ISRCTN	ISRCTN wordt niet meer aangevraagd

## Resultaten

### Samenvatting resultaten

1. Debats IB, Booij DI, Deutz NE, Buurman WA, Boeckx WD, van der Hulst RR. Infected chronic wounds show different local and systemic arginine conversion compared with acute wounds. *J Surg Res.* 2006 Aug;134(2):205-14.
2. Booij DI, Debats IB, Boeckx WD, van der Hulst RR. Risk factors and blood flow in the free TRAM flap: Smoking and high flap weight impair the free TRAM flap microcirculation. *Ann Plast Surg.* 2007 Oct; 59(4):364-71
3. Booij DI, Debats IB, Boeckx WD, van der Hulst RR. Laser Doppler Flowmetry in the free TRAM flap when using zone IV: a clinical study indicating a 48 hour delay in choke vessel opening. *JPRAS* 2008 March; 61(3):282-8
4. Debats IB, Booij DI, Wehrens KWE, Van den Hogen E, Deutz NE, Bemelmans MHA, van der Hulst RR. Oral arginine supplementation and the effect of skin graft donor sites; a randomized clinical pilot study. Accepted for publication in *J Burn Care and Res.*
5. Booij DI, Wehrens KWE, Lievaart V, Debats IB, Marcus MAE, van der Hulst RR. Peri-operative fluid overload increases postoperative complications in the free TRAM flap. Submitted for publication.