

# Effect of high frequency rTMS on negative symptoms and cognitive functioning in schizophrenia

Gepubliceerd: 01-03-2008 Laatste bijgewerkt: 18-08-2022

Brain activity in people with schizophrenia differs when compared to brain activity of people who do not have this condition. There is evidence that the negative symptoms of schizophrenia are related to reduced cortical activation involving the...

<b>Ethische beoordeling</b>	Positief advies
<b>Status</b>	Werving nog niet gestart
<b>Type aandoening</b>	-
<b>Onderzoekstype</b>	Interventie onderzoek

## Samenvatting

### ID

NL-OMON25191

### Bron

Nationaal Trial Register

### Verkorte titel

TreNSS

### Aandoening

- schizophrenia
- negative symptoms
- transcranial magnetic stimulation
- TMS
- cognitive functioning

### Ondersteuning

**Primaire sponsor:** University Medical Center Groningen (UMCG)

### Onderzoeksproduct en/of interventie

# Uitkomstmaten

## Primaire uitkomstmaten

The primary outcome of the study is a significant decline of negative symptoms and cognitive dysfunctioning after bilateral high frequency rTMS treatment of the DLPFC.

## Toelichting onderzoek

### Achtergrond van het onderzoek

Negative symptoms in schizophrenia include anhedonia, avolition, alogia and blunted affect. They predict a worse clinical outcome and are often indicative of poorer social, occupational and global outcomes. Currently, no effective treatment exists to improve negative symptoms. The primary objective of the current investigation is to determine the effectiveness of bilateral high frequency rTMS treatment of the dorsolateral prefrontal cortex (DLPFC) on negative symptoms and cognitive dysfunction in patients with schizophrenia. The secondary objective is to study brain activity and changes in brain activity after rTMS through a fMRI study.

Design: Double-blind sham controlled study.

Sample size: 32 patients, 16 in the treatment group and 16 in the sham group.

Study population: All patients should be over 18 years old and should meet the diagnostic criteria for schizophrenia, based on the patient's medical file and confirmed by a standardized psychiatric assessment. Inpatients as well as outpatients can participate.

Intervention: The rTMS group of 16 patients will receive bilateral rTMS stimulation during 15 days, sessions being given twice daily. The sham group of 16 patients will constitute the placebo group and receive 15 days of bilateral sham stimulation twice daily. The primary outcome of the study is a significant decline of negative symptoms and cognitive dysfunctioning after bilateral high frequency rTMS treatment of the DLPFC. The secondary outcome of the study is increased cortical activation in the DLPFC after rTMS treatment, measured through a fMRI study.

### Doel van het onderzoek

Brain activity in people with schizophrenia differs when compared to brain activity of people who do not have this condition. There is evidence that the negative symptoms of schizophrenia are related to reduced cortical activation involving the prefrontal cortex, in particular the activation in the dorsolateral prefrontal cortex (DLPFC) seems to be reduced. The reduced activity in the frontal cortex correlates with the severity of negative symptoms. Furthermore, several publications on functional neuroimaging literature suggest that

hypofrontality in schizophrenia may be related to both negative symptoms and cognitive deficits. Transcranial magnetic stimulation (TMS) is a non-invasive, save method to stimulate the cerebral cortex and thereby alter neuronal function. In TMS a coil, which delivers brief magnetic pulses is placed adjacent to the scalp. The magnetic field passes the skull without hindrance and induces an electric current in certain brain regions. The neuronal pathways may be excited or inhibited, depending on the intensity and frequency of stimulation. Low frequency TMS (1 Hz) decreases brain activity and high frequency TMS (>5 Hz) increases brain activity.

As mentioned earlier, the brain activity in the DLPFC is reduced in people suffering from schizophrenia. Increasing brain activity in the DLPFC by using bilateral high frequency rTMS might therefore prove an effective treatment of negative symptoms and cognitive functioning in schizophrenia.

### **Onderzoeksopzet**

Follow-up measurements will be done at 4 weeks and 3 months

### **Onderzoeksproduct en/of interventie**

The rTMS group of 16 patients will receive bilateral high frequency rTMS stimulation during 15 days, sessions being given twice daily.

The sham group of 16 patients will constitute the placebo group and receive 15 days of bilateral sham stimulation twice daily.

## **Contactpersonen**

### **Publiek**

J. Dlabac-de Lange  
Groningen  
The Netherlands

### **Wetenschappelijk**

J. Dlabac-de Lange  
Groningen  
The Netherlands

## **Deelname eisen**

## **Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)**

1. Participants must be 18 years or older.
2. They must be diagnosed with schizophrenia and the diagnosis must be confirmed by a standardised psychiatric assessment, the Schedules for Clinical Assessment in Neuropsychiatry (SCAN 2.1).
3. Patients included must have prominent negative symptoms and patients must have a negative subscore = or > than 15 on the Positive and Negative Syndrome Scale for Schizophrenia (PANSS).
4. Only patients who are fully capable of making their own decision regarding participation in the research will be included.

## **Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)**

1. Exclusion criteria include rTMS and MRI contraindications (e.g. a personal or family history of epileptic seizures, history of brain surgery, intracerebral or pacemaker implants, inner ear prosthesis or other metal prosthetics/implants).
2. Neurological disorders, a head injury with loss of consciousness in the past.
3. Substance dependency within the previous 6 months.
4. Previous treatment with rTMS.
5. Severe behavioural disorders.
6. Inability to provide informed consent.
7. Claustrophobia.
8. Female patients who are pregnant will be excluded.

## **Onderzoeksopzet**

## Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Dubbelblind
Controle:	Placebo

## Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-05-2008
Aantal proefpersonen:	32
Type:	Verwachte startdatum

## Ethische beoordeling

Positief advies	
Datum:	01-03-2008
Soort:	Eerste indiening

## Registraties

### Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

### Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

### In overige registers

Register	ID
NTR-new	NL1216

**Register**

NTR-old

Ander register

ISRCTN

**ID**

NTR1261

METC : 2008/058

ISRCTN wordt niet meer aangevraagd

## Resultaten

**Samenvatting resultaten**

N/A