

Improving medication adherence with Treatment Adherence Therapy.

Gepubliceerd: 07-12-2011 Laatste bijgewerkt: 18-08-2022

Compared to treatment as usual adding a tailored adherence interventions Treatment Adherence Therapy (TAT) for outpatient with schizophrenia will improve medication adherence.

Ethische beoordeling	Niet van toepassing
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON25257

Bron

Nationaal Trial Register

Verkorte titel

none

Aandoening

Patients with schizophrenia and poor medication adherence

Ondersteuning

Primaire sponsor: Arkin Amsterdam

Overige ondersteuning: Arkin,
Stichting Agis

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The proportion of non-adherent patients based on the Brief Adherence Rating Scale (BARS).

Toelichting onderzoek

Achtergrond van het onderzoek

Background:

It is well established that approximately 50% of patients with schizophrenia have poor medication adherence. In general this results in increased risk of relapse and hospitalisation, and poorer outcomes. Up to date there is no adherence intervention with sufficient evidence base for its efficacy. Adherence interventions usually target one of a variety of reasons for non-adherence. With TAT it is possible to tailor the adherence intervention to the reasons of non-adherence. A first study of the effectiveness of TAT has shown that medication adherence significantly improved (Starring et al 2010; NTR1159).

Research Question:

In this randomised controlled trial we want to examine the effectiveness of TAT on clinical outcomes in patients with schizophrenia.

Methode/design:

This is a randomised controlled trial. In total 745 outpatients with schizophrenia will be screened for medication adherence. Patients with poor medication adherence, and who give informed consent will be included. Included patients will be allocated to the control condition or the experimental condition. In the control condition, patients receive care as usual according to the principles of FACT. In the experimental condition, patients receive care as usual and in addition will be offered TAT performed by a trained psychologist who is not the regular key worker of the patient. Measurements will be performed at baseline, and after 12 (T1), and 24 months (T2).

Sample size calculation/data analysis:

At baseline all patients will be non-adherent. A large scale study amongst 34.128 patients with schizophrenia showed that 30% of NA patients will be adherent the following year (35). This corresponds with a study based on 2% of the Dutch population which showed that 23% of NA patients (using antipsychotics) were only temporarily NA (36). In this study we expect 70% of NA patients to remain non-adherent in the control condition after 12 months. We expect this to be 50% in the experimental condition. This is considered a clinically meaningful effect of MI. To detect this difference in a 1-sided test ($\alpha=0.05$; $1-\beta=0.80$) we need

150 participants in total. Analyses are according intention to treat principles.

Doel van het onderzoek

Compared to treatment as usual adding a tailored adherence interventions Treatment Adherence Therapy (TAT) for outpatient with schizophrenia will improve medication adherence.

Onderzoeksopzet

Baseline measurements will be conducted after randomization and follow-up measurements will be conducted at 12, and 24 months after the baseline measurement.

Onderzoeksproduct en/of interventie

Treatment as usual:

Treatment as usual consists of regular outpatient care according to the principles of FACT.

TAT:

Treatment Adherence Therapy (TAT) has recently been developed by Prof Dr Niels Mulder en Dr Tonnie Staring (Erasmus Universiteit, Rotterdam). TAT consists of three main therapeutical modules, based on an empirical theoretical model developed by Staring (2006). Modules will be deployed based on an extensive assessment of the underlying cause of non adherence. TAT is therefore tailored to needs and situation of the patient. TAT consists of 10 individual weekly sessions and 3 monthly individual booster sessions.

Contactpersonen

Publiek

Arkin Academy

Klaprozenweg 111
M.J. Kikkert
Amsterdam 1033 NN
The Netherlands
+31 (0)20 5904422

Wetenschappelijk

Arkin Academy

Klaprozenweg 111
M.J. Kikkert
Amsterdam 1033 NN
The Netherlands
+31 (0)20 5904422

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Diagnosis schizophrenia according to DSM-IV criteria;
2. Current prescription of an antipsychotic medication;
3. Outpatient and autonomous in collecting and using his or her antipsychotic medication;
4. Poor medication adherence.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Unable to follow TAT treatment due to inadequate mastery of the Dutch language, or severe cognitive impairment;
2. Severe substance abuse.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd

Blinding:	Open / niet geblindeerd
Controle:	Actieve controle groep

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-03-2012
Aantal proefpersonen:	150
Type:	Verwachte startdatum

Ethische beoordeling

Niet van toepassing	
Soort:	Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL3035
NTR-old	NTR3183
Ander register	:
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A