

Nerve Growth Factor & validation study

Gepubliceerd: 30-01-2018 Laatst bijgewerkt: 15-05-2024

Dried blood spot analysis is a reliable method to measure Nerve Growth Factor &.

Ethische beoordeling	Positief advies
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Observationeel onderzoek, zonder invasieve metingen

Samenvatting

ID

NL-OMON25927

Bron

NTR

Aandoening

Nerve growth factor β

Healthy volunteers

Ondersteuning

Primaire sponsor: VU University Medical Center, Amsterdam

Overige ondersteuning: Sponsor

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The main study parameters will be the correlation between the DBS and saliva methods compared to venous blood sampling. In order to do so, NGF-& concentrations (pg/ml) for each method will be measured. The difference between saliva testing and venous blood and the difference between dried blood spot and venous blood will be calculated.

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale: There are roughly two types of ovulation: spontaneous and induced. Copulation is the speculated mechanism behind induced ovulation. The ovulation induction substance in semen has been identified as nerve growth factor β (NGF- β). In animal studies, semen plasma derived NGF- β indeed has an effect on female ovulation, follicle development and a luteotropic effect. The role of NGF- β on human ovulation has never been studied. It is known that NGF- β plays an important role in many different types of human tissue, for instance the nervous system and immune system. All studies use venous blood to measure the NGF- β concentration. However, in order to study the effect of NGF- β on human ovulation, multiple serial measurements are needed, ideally immediately before and after natural intercourse. Therefore, a patient-friendly and minimally interrupting method is necessary. Serial dried blood spot analysis and saliva sample testing are potential methods, but both are not validated for NGF- β yet. The aim of this study will be to do a method comparison study, to validate the use of both methods compared to venous blood for NGF- β .

Objective: To answer the question if dried blood spot (DBS) analysis and saliva testing are reliable methods to determine the concentration of NGF- β in healthy subjects.

Study design: Cross sectional method comparison validation study.

Study population: Healthy human volunteers of reproductive age (18 - 43 years).

Main study parameters/endpoints: The main study parameters are the correlation and limits of agreement between dried blood spots/saliva samples compared to venous blood concentration.

Nature and extent of the burden and risks associated with participation, benefit and group relatedness: The procedure for participants includes one hospital visit or home visit for saliva, dried blood spot and venous blood sampling and a questionnaire has to be completed. There will be no extra risks nor benefits for the participants. The risk analysis for this study concluded the risk for (serious) adverse events is negligible.

Doe~~l~~ van het onderzoek

Dried blood spot analysis is a reliable method to measure Nerve Growth Factor β .

Onderzoeksopzet

Questionnaire will be completed at home, before the first visit. During the first visit, all samples will be collected. Afterwards all samples will be analyzed.

Onderzoeksproduct en/of interventie

Questionnaire, saliva sampling, serum sampling, dried blood spot collection.

Contactpersonen

Publiek

VUMC, Department of Obstetrics and Gynaecology / Division of Reproductive Medicine

Nienke Schouten
De Boelelaan 1118 (PK 0 Z 116.1)

Amsterdam 1081 HZ
The Netherlands
+312044442867

Wetenschappelijk

VUMC, Department of Obstetrics and Gynaecology / Division of Reproductive Medicine

Nienke Schouten
De Boelelaan 1118 (PK 0 Z 116.1)

Amsterdam 1081 HZ
The Netherlands
+312044442867

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

- Both male and female volunteers.
- 18 years or older.
- Willingness to give informed consent, to donate venous blood, dried blood spots obtained through a finger puncture, saliva and complete a questionnaire.
- Sufficient command of Dutch or English language and capable of understanding participants information.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- Contraindications for donating blood (phlebitis, dermatitis, psoriasis, lymphedema, arterial venous fistula, hematoma on or around insert place, mamma-amputation or axillary dissection of lymph nodes).
- Age younger than 18 years.
- Poor dental condition.
- Volunteers who brushed their teeth within 45 minutes prior to sample collection.
- Volunteers who had dental work within 24 hours prior to sample collection.
- Volunteers who had their last meal, cigarette, or drank anything except for water within 60 minutes prior to sample collection.

Onderzoeksopzet

Opzet

Type:	Observationeel onderzoek, zonder invasieve metingen
Onderzoeksmodel:	Anders
Toewijzing:	N.v.t. / één studie arm
Blinding:	Open / niet geblindeerd

Controle: N.v.t. / onbekend

Deelname

Nederland
Status: Werving nog niet gestart
(Verwachte) startdatum: 07-02-2018
Aantal proefpersonen: 60
Type: Verwachte startdatum

Ethische beoordeling

Positief advies
Datum: 30-01-2018
Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 44502
Bron: ToetsingOnline
Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL6804
NTR-old	NTR6990
CCMO	NL63048.029.17
OMON	NL-OMON44502

Resultaten