

Initial non-operative treatment versus direct operative treatment for simple appendicitis in children.

Gepubliceerd: 20-07-2016 Laatste bijgewerkt: 18-08-2022

Initial non-operative treatment strategy is associated with as many or fewer complications, a better health-related QOL, reduced costs, and it avoids surgery in the majority of patients.

Ethische beoordeling	Niet van toepassing
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON26428

Bron

Nationaal Trial Register

Verkorte titel

APAC study

Aandoening

Appendicitis

Ondersteuning

Primaire sponsor: Academisch medisch centrum Amsterdam

Overige ondersteuning: This study is funded by ZONmw

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The primary outcome is the proportion of patients experiencing complications within one

month follow-up. An adjudication committee will be installed to review the complications. They will review all patients with possible complications to determine whether or not they fulfill the criteria for complications. All Complications will be recorded.

Complications are defined as:

- Allergic reaction to antibiotics administered. In case an allergic reaction is suspected, the child will be referred to the allergist for further evaluation.
- Re-admission for an indication other than recurrent appendicitis but related to appendicitis (such as readmission for observation of fever or abdominal pain)

Complications associated with appendectomy:

- Superficial Site infection
- Intra-abdominal abscess
- Stump leakage/stump appendicitis
- Secondary / prolonged Bowel Obstruction
- Anesthesia Related complications (such as pneumonia)
- Hernia cicatricialis
- Need for other surgical or radiological intervention than appendectomy but related to appendicitis (such as percutaneous drainage of an abscess, surgical intervention for a superficial site infection)

Toelichting onderzoek

Doel van het onderzoek

Initial non-operative treatment strategy is associated with as many or fewer complications, a better health-related QOL, reduced costs, and it avoids surgery in the majority of patients.

Onderzoeksopzet

Outcomes will be measured at discharge, 7 days, 1, 6, and 12 months after randomization as described above.

Onderzoeksproduct en/of interventie

Intervention group (Non-operative treatment strategy): Clinical observation for 48 hours with administration of Intravenous administration of amoxicillin/clavulanic acid 25/2.5mg 6-hourly (total 100/10 mg/kg daily; maximum 6000/600mg a day) and gentamicin 7mg/kg once daily for 48 hours. If after 48 hours the patient fulfils the predefined discharge criteria, the antibiotics will be switched to oral amoxicillin/clavulanic acid 50/12.5 mg/kg 8-hourly (max 1500/375mg a day) for in total 7 days and discharge. An appendectomy is reserved for those patients with clinical deterioration, non-improvement after 72 hours or recurrent appendicitis.

Control group (Operative treatment strategy): Clinical observation and semi-urgent appendectomy. Pre-, peri- and postoperative care according to local protocol. No routine postoperative antibiotics. Discharge if the patient fulfils the predefined discharge criteria.

Contactpersonen

Publiek

Kinderchirurgisch Centrum Amsterdam, Emma's Kinderziekenhuis AMC & VU medisch centrum
R.R. Gorter
Amsterdam
The Netherlands
+31 (0)20 4444444

Wetenschappelijk

Kinderchirurgisch Centrum Amsterdam, Emma's Kinderziekenhuis AMC & VU medisch centrum
R.R. Gorter
Amsterdam
The Netherlands
+31 (0)20 4444444

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Inclusion criteria:

Eligible for inclusion are all children from 7 to 17 years old, inclusive, with a radiologically confirmed simple appendicitis.

Definition of simple appendicitis is based upon predefined clinical, biochemical and radiological (ultrasound) criteria.

Clinical & biochemical criteria:

- Unwell but not generally ill

- Localized tenderness in the right iliac fossa region
- Normal/hyperactive bowel sounds
- No guarding or palpable mass
- Biochemical signs of infection (Elevated White Blood Cell count (WBC) and/or C-reactive protein (CRP)).

As recommended by the national guideline, all children with a clinical and/or biochemical suspicion should undergo ultrasound studies. Ultrasound criteria to confirm the diagnosis of acute simple appendicitis are:

- An incompressible, painful appendix with an outer diameter > 6 mm
- Secondary signs of inflammation such as surrounding fat infiltration, limited clear free fluid surrounding the appendix, hyperemia within the appendiceal wall.
- No fecolith, no signs of perforation, no signs of intra-abdominal abscess or phlegmone.

In case the ultrasound is inconclusive, additional imaging studies may be obtained. CT-scan is not recommended in the young children (due to its risk of radiation induced malignancy). MRI is recommended in those places with sufficient experience in the interpretation of the results. Only those in whom imaging studies confirm the diagnosis of simple appendicitis can be included.

In case there is no certain diagnosis and a “watchful waiting” strategy is chosen, the patient cannot be included.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- Generalized peritonitis, complex appendicitis or sepsis (based upon predefined criteria and scoring system).

Scoring system: A scoring system was developed determining the risk of complex appendicitis based upon five pre-operative variables. Points have been awarded to each variable. In case the total score is less than 4 points, the patient is likely to have a simple appendicitis. In case the score is 4 or more points, the chance of having complex appendicitis is

significant and those children will be excluded from this study.

Variables:

Diffuse abdominal guarding (3 points)

CRP level more than 38 mg/L (2 points)

Signs on ultrasound indicative of complex appendicitis (2 points)

More than one day abdominal pain (2 points)

Temperature: more than 37.5 degree Celsius (1 point)

- Fecolith (ultrasound)
- Serious co-morbidity
- Recurrent appendicitis
- Suspicion of an underlying malignancy or inflammatory bowel disease
- Documented type 1 allergy to the antibiotics used.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	Actieve controle groep

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-12-2016
Aantal proefpersonen:	334
Type:	Verwachte startdatum

Ethische beoordeling

Niet van toepassing

Soort:

Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL5822
NTR-old	NTR5977
Ander register	: APAC2016

Resultaten