

A phase II study of up-front red blood cell transfusion followed by maintenance Erythropoetin-alpha (Epo-alpha) s.c. support during chemotherapy of solid tumors.

Gepubliceerd: 19-09-2005 Laatste bijgewerkt: 18-08-2022

An upfront RBCT aiming at low-normal Hb levels will ameliorate anemia-caused tumor hypoxia-related resistance to chemotherapy before the start of chemotherapy and may decrease secondary anemia-induced endogenous release of cytokines like VEGF,...

Ethische beoordeling	Positief advies
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON26479

Bron

NTR

Verkorte titel

pCATS

Aandoening

Solid tumors

Ondersteuning

Primaire sponsor: Not applicable.

Overige ondersteuning: Ortho Biotech / Jansen-Cilag B.V., the Comprehensive Cancer Center West (Integraal Kankercentrum West)

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Hb levels before the start of and during chemotherapy; safety of the pCATS anemia treatment regimen.

Toelichting onderzoek

Achtergrond van het onderzoek

Supportive care treatment of anemia with a blood transfusion before the start of standard chemotherapy, maintained with weekly erythropoietin-alpha subcutaneously injections during the duration of the chemotherapy

Doel van het onderzoek

An upfront RBCT aiming at low-normal Hb levels will ameliorate anemia-caused tumor hypoxia-related resistance to chemotherapy before the start of chemotherapy and may decrease secondary anemia-induced endogenous release of cytokines like VEGF, osteopontin. The maintenance of optimal Hb levels at this lower-normal range during chemotherapy by weekly maintenance administration of Epo-alpha s.c. at doses with proven safety and efficacy creates optimal conditions for tumor oxygenation, without the presumably high-Hb level associated adverse effects.

Onderzoeksproduct en/of interventie

Anemia-treatment consisting of preventive RBCT before the start of chemotherapy followed by the maintenance administration of Epo-alpha s.c. during chemotherapy.

Contactpersonen

Publiek

't Lange Land Ziekenhuis, Department of Internal Medicine,
P.O. Box 3015
A.C. Ogilvie
Zoetermeer 2700 KJ

The Netherlands
+31 (0)79 3462881

Wetenschappelijk

't Lange Land Ziekenhuis, Department of Internal Medicine,
P.O. Box 3015
A.C. Ogilvie
Zoetermeer 2700 KJ
The Netherlands
+31 (0)79 3462881

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Histological or cytological documentation of solid tumor (breast- or colorectal- or ovarian- or lung- or esophageal- or stomach- or bladder- or prostate- or germ cell- or cervical cancer or sarcoma)
2. Age ≥ 18 years;
3. ECOG performance status of 0, 1 or 2;
4. Being scheduled to receive chemotherapy or having received already 1 cycle of chemotherapy and being scheduled to receive at least 3 cycle of chemotherapy prior to study entry;
5. Life expectancy of at least 6 month;
6. Signed written informed consent obtained prior to study entry;
7. Anemia: Hb < 7.0 mmol/L tested within 7 days before enrolment;
8. Adequate bone marrow function as assessed within 7 days before enrolment by:
 - a. Absolute neutrophil count $\geq 1.5 \times 10^9/L$;
 - b. Platelets $\geq 100 \times 10^9/L$;
9. Iron status measurements including levels of ferritin, transferrin, iron and iron saturation within 7 days after enrolment;
10. Patient is able to comply with scheduled follow up.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Excluded medical conditions:

1. Having more than 1 cycle of the current chemotherapy administered prior to inclusion;
2. Having 1 cycle of chemotherapy administered before inclusion and scheduled to receive less than 3 additional cycles;

3. Untreated folate or cobalamin deficiency;
4. Untreated haemolytic anemia defined by decreased serum haptoglobin levels;
5. Anemia due to hypoproliferative or maturation bone marrow disorders;
6. Clinically evident untreated congestive heart failure;
7. Serious, untreated cardiac arrhythmias;
8. Symptoms of untreated coronary heart disease or ischemia;
9. Untreated hypertension;
10. History of HIV infection.

Excluded therapies, medications and conditions, previous and concomitant:

11. Androgen treatment within 2 month before enrolment;
12. Anti-cancer chemotherapy or immunotherapy within 4 weeks of study entry;
13. Darbepoetin or erythropoetin treatment within 4 weeks before enrolment;
14. Bone marrow transplantation or stem cell transplantation within 4 months of study entry;
15. Investigational drug therapy within 4 weeks of study entry or during this study,
16. Pregnant or breast-feeding patients. Women of childbearing potential must have a negative pregnancy test performed within 7 days of the start of treatment. Adequate birth control measures will be required during the course of the trial,
17. Known or suspected allergy to Epo-alpha.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Anders
Blinding:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving gestopt
(Verwachte) startdatum:	15-09-2005
Aantal proefpersonen:	23
Type:	Werkelijke startdatum

Ethische beoordeling

Positief advies

Datum: 19-09-2005
Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL369
NTR-old	NTR409
Ander register	: N/A
ISRCTN	ISRCTN81146641

Resultaten

Samenvatting resultaten

N/A