

PTSD as a result of chronic interpersonal violence in early childhood; Imaginal Exposure vs Imagery Rescripting vs Body Focused Rescripting.

Gepubliceerd: 13-02-2013 Laatste bijgewerkt: 18-08-2022

Objective of the study: The research aims to 2 new forms of treatment for patients with comorbid PTSD as a result of multiple and repeated abuse in childhood to assess and compare Imaginary Exposure. The expectation is that these new forms of...

Ethische beoordeling	Positief advies
Status	Werving tijdelijk gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON26598

Bron

Nationaal Trial Register

Verkorte titel

PTSD in early childhood

Aandoening

The research aims to 2 new forms of treatment for patients with comorbid PTSD as a result of multiple and repeated abuse in childhood to assess and compare Imaginary Exposure. The expectation is that these new forms of treatment will lead to a better end-state functioning, less dropout and a higher valuation for feasibility of therapists. The expectation is that Body Focused Rescripting superior to Imaginary Rescripting.

Hypotheses treatment study (1): Effects 1. We expect treatment effects on the primary outcome measures in the ranking IE

Ondersteuning

Primaire sponsor: PSYQ Haaglanden

Carel Reinierszkade 197

2593 HR Den Haag

Nederland

Tel: 088 3573900

Email: psychotrauma-onderzoek@psyq.nl

Overige ondersteuning: PSYQ (Parnassia groep) is the main sponsor of this study.

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

1. Anger Expression Scale (AEQ);

2. State-Trait Anger Inventory (ZAV);

3. Guilt scale (Kubany);

4. Shame scale (Smucker);

5. Grief / consolation / happiness scale.

Toelichting onderzoek

Achtergrond van het onderzoek

Meta-analyzes show that Trauma-Focused Cognitive Behavioral, Imaginary Exposure and Eye Movement

Desensitization-Reprocessing the most effective psychological treatments for posttraumatic stress disorder (PTSD) (Bisson et al, 2007; Bradley, Greene, Russ, Dutra & West, 2005; Cloitre, 2009; Seidler & Wagner, 2006). In 40-70% of the patients indicate that relatively short treatment (9-12 sessions of 90 minutes) to considerable reduction of PTSD symptoms. The vast majority of research has been conducted on PTSD following following a single traumatic event in childhood .

There is still little research on the application of this protocols to PTSD as a result of repeated and prolonged interpersonal

abuse in childhood (eg sexual and / or physical abuse) within the immediate environment.

The question arises whether "new" forms of treatment such as Imaginary Rescripting and Body Focused Rescripting that more active intervention on key processes that play a role in the development of PTSD as dysfunctional based schedules and tonic immobility (TI) effective a proven effective form of treatment as imaginal exposure (IE). In this study, therefore the effectiveness of three treatments for chronic PTSD after interpersonal trauma in childhood investigated, namely (1) imaginal exposure, (2) Imaginary Rescripting and (3) Body Focused Rescripting.

Doel van het onderzoek

Objective of the study:

The research aims to 2 new forms of treatment for patients with comorbid PTSD as a result of multiple and repeated abuse in childhood to assess and compare Imaginary Exposure. The expectation is that these new forms of treatment will lead to a better end-state functioning, less dropout and a higher valuation for feasibility of therapists. The expectation is that Body Focused Rescripting superior to Imaginary Rescripting.

Hypotheses treatment study (1):

Effects 1. We expect treatment effects on the primary outcome measures in the ranking IE
A. In Rescripting therapies is influenced not only the fear but also anger, guilt, shame and sadness;

B. Because context dependence is less (UCS revaluation);

C. Because active intervention is essential processes in development of PTSD, such as TI;

D. Body Focused Rescripting better effects than ImRes because ImRes will initially focus on the explicit memory, while patients with TI and dissociation possibly better results expected if one focuses on implicit memory. The context of the UCS / UCR would be better represented if the traumatic event is not in vitro (imaginary) but in vivo depicts. The newly learned response to the UCS representation is not directly aimed at expressing the blocked emotion (emotional processing), but had to be aware of sensorimotor components (sensorimotor processing) such as perception of sensory and physiological sensations of fixed action tendencies and defensive postures focus more on implicit than explicit memory (Ogden, Minton & Pain, 2006).

2. Dropout. We expect that the IE condition the dropout % higher than in the latter treatments because BFRes ImRes and less stress.

3. Preferences of therapists. It is expected that therapists 'new' treatments find more workable and less heavy for the patient seem to be. Hypothesized predictors of treatment success (2): There will be investigated predictors of treatment success. The hypotheses to be tested, we formulate based on the most recent empirical findings. If predictors are included: severity axis II (borderline symptoms and avoidant symptoms), TI (TI-self-report and TI as discrepancy between low physiological and subjective anxiety), dissociation, alcohol and drug use, severity of trauma, therapeutic relationship and HRV . We expect that

patients with high TI and high dissociation score a better effect on the ImRes condition and even more in the BFRes condition than the IE condition. In the patients with a low dissociation TI and low score will be the difference between the conditions are less severe.

Generates motor (tonic) immobility post-traumatic complications? Serious sexual abuse involving penetration increases the likelihood of experiencing motor (tonic) immobility (Heidt et al., 2005)? Is motor (tonic) immobility one aspect of peritraumatic dissociation (Hagenaars, et al, 2009)? What correlation is there between motor (tonic) immobility, peritraumatic dissociation, PTSD, anxiety and depressive (Heidt et al., 2005)?

Onderzoeksopzet

The waiting period of 5 weeks of ring forms include the control group. Patients are randomized to 1 of 3 treatments assigned.

Two sessions of 90 minutes per week for 6 weeks last 4 weeks 1 time per week. Total 16 seats plus Homework 2 times per week (max. 1 hour).

Duration of treatment 11 weeks.

Homework: • IE / ImRes / BF Res: listening to the tape of the session 2 times a week Exposure in vivo is in all conditions outside the study protocol. 52 patients per condition.

Measurement 1 after inclusion, measuring 2 to start treatment after 5 weeks of Measurement 3, Measurement 4 after

10 weeks (termination therapy) and follow-up after 3 months (Measure 5) and 12 months (Measure 6). In all conditions

of the research will end a break of 3 months be inserted after assessing whether the therapy according to TAU

(treatment as usual) should be prosecuted. During the pause, the handler if necessary by phone. As further

improvements after termination of the protocol are expected, a pause of 3 months at comparable RCTs are necessary.

Onderzoeksproduct en/of interventie

In this study the effectiveness of three treatments for chronic PTSD after interpersonal trauma in childhood will be investigated, namely (1) imaginal exposure (IE), (2) Imaginary Rescripting (ImRes) and (3) Body Focused Rescripting (BFR es).

Study design:

This study is a multicenter randomized 3-group trial.

3 conditions: IE, ImRes, BFRes. The waiting period of 5 weeks of ring forms include the control group. Patients are randomized to 1 of 3 treatments assigned. Two sessions of 90 minutes per week for 6 weeks last 4 weeks 1 time per week. Total 16 seats plus homework 2 times per week (max. 1 hour).

Duration of treatment 11 weeks.

Homework: IE / ImRes / BF Res: listening to the tape of the session 2 times a week. Exposure in vivo is in all conditions outside the study protocol.

52 patients per condition.

Measurement 1 after inclusion, measuring 2 to start treatment after 5 weeks of Measurement 3, Measurement 4 after 10 weeks (termination therapy) and follow-up after 3 months (Measure 5) and 12 months (Measure 6).

In all conditions of the research will end a break of 3 months be inserted after assessing whether the therapy according to TAU (treatment as usual) should be prosecuted. During the pause, the handler if necessary by phone. As further improvements after termination of the protocol are expected, a pause of 3 months at comparable RCTs are necessary.

Background of the study:

Meta-analyzes show that Trauma-Focused Cognitive Behavioral, Imaginary Exposure and Eye Movement

Desensitization-Reprocessing the most effective psychological treatments for posttraumatic stress disorder (PTSD) (Bisson et al, 2007; Bradley, Greene, Russ, Dutra & West, 2005; Cloitre, 2009; Seidler & Wagner, 2006). In 40-70% of the patients indicate that relatively short treatment (9-12 sessions of 90 minutes) to considerable reduction of PTSD symptoms. The vast majority of research has been conducted on PTSD following following a single traumatic event in childhood .

There is still little research on the application of this protocols to PTSD as a result of repeated and prolonged interpersonal abuse in childhood (eg sexual and / or physical abuse) within the immediate environment. The question arises whether "new" forms of treatment such as Imaginary Rescripting and Body Focused Rescripting that more active intervention on key processes that play a role in the development of PTSD as dysfunctional based schedules and tonic immobility (TI) effective a proven effective form of treatment as imaginal exposure (IE). In this study, therefore the effectiveness of three treatments for chronic PTSD after interpersonal trauma in childhood investigated, namely (1) imaginal exposure, (2)

Contactpersonen

Publiek

P.M. Jong, de
Psychotherapy & Moment
drs. P.M. de Jong & M.T.C van Hemert
Sarphatistraat 27a
Amsterdam 1018 EV
The Netherlands
0031 (0) 20- 528 52 31

Wetenschappelijk

P.M. Jong, de
Psychotherapy & Moment
drs. P.M. de Jong & M.T.C van Hemert
Sarphatistraat 27a
Amsterdam 1018 EV
The Netherlands
0031 (0) 20- 528 52 31

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

PTSD patients from mainstream mental health. The sample consists of 156 patients with a primary diagnosis of PTSD and comorbid disorders reported to various treatment programs PsyQ Psychotrauma and Personality Problems The Hague, Rotterdam and Spijkenisse. The various Psychotrauma treatment programs are mainly patients with a primary diagnosis of PTSD, is also expected that around 50% of patients at intake to meet the inclusion criteria (total inflow Indoor 1, 5 years). Participants are victims of repeated or chronic interpersonal trauma in childhood (eg sexual or physical abuse) at an age younger than 16 years. Participants are 18 years or older.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Psychiatric problems that may interfere with the study participation or that require more intensive care than can be offered in the present study , including dementia, psychotic symptoms, depression with suicidal ideation, full blown borderline personality disorder, substance dependence, dissociative identity disorder IV no fixed residence, major financial problems, no aid figure, problems with police and law, current abuse;
2. Current use of tranquillizers;
3. On as IV no fixed residence, major financial problems, no aid figure, problems with police and law, current abuse.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	Actieve controle groep

Deelname

Nederland	
Status:	Werving tijdelijk gestopt
(Verwachte) startdatum:	01-03-2013
Aantal proefpersonen:	156
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	13-02-2013

Soort:

Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL3702
NTR-old	NTR3872
CCMO	NL42151.018.12
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

Not yet. The first general publication will be coming in april.