

A multicenter, double blind placebo controlled randomized trial for Benign esophageal Anastomotic Strictures: Savary dilation vs savary dilation with TriAmcinolon.

Gepubliceerd: 09-03-2010 Laatste bijgewerkt: 18-08-2022

By injection of triamcinolone before dilation we hypothesise that the time to repeat dilation is increased and the total number of dilation sessions is decreased as well.

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON26668

Bron

Nationaal Trial Register

Verkorte titel

BASTA-study

Aandoening

Patients with dysphagia after esophagectomy with cervical anastomosis due to a benign anastomotic stricture.

Patienten met dysfagie na een buismaagreconstructie na slokdarmresectie met een cervicale anastomose veroorzaakt door een benigne anastomotische strictuur

Ondersteuning

Primaire sponsor: Maag-, darm-, leverziekten UMC Utrecht

Overige ondersteuning: UMC Utrecht (investigator driven)

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Number of patients dysphagia free after 1 dilation in a period of 6 months (success rate).

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale: Dysphagia due to benign esophageal stricture formation can significantly impair quality of life. Stricture formation at the site of the anastomosis after esophagectomy is increasingly reported as a common cause of benign esophageal strictures, whereas reflux disease less frequently causes stricture formation, probably due to the increased use of PPI's (proton pump inhibition).

Strictures can be classified according to complexity. In general, peptic strictures are simple, focal and straight and require only 1-3 dilations to relieve dysphagia. Complex strictures are longer (>2 cm), angulated, irregular or with a severely narrowed diameter. The most common complex strictures include post-radiation, caustic or anastomotic strictures. These strictures are frequently refractory to dilation and require multiple (>5) sessions sometimes at weekly intervals.

Adequate treatment of these strictures is of utmost importance. Dilation is the mainstay of this treatment and can be performed by balloon dilatation or by Savary Gilliard bougienage. Both treatments have shown to be equally effective and safe, however Savary Gilliard dilations are re-useable, making this a more cost-effective method.

Addition of intra-esophageal triamcinolon injections to dilatation was first mentioned in 1969, but this technique has only been increasingly employed over the last decade. Several prospective studies have demonstrated its use in dogs, adults and children. Triamcinolon was found to be safe and effective in lengthening the dilation-free interval and reducing the risk for recurrent stricture formation in patients with strictures of all causes, but also in prospective series with peptic or corrosive strictures solely. However, a major disadvantage of these studies is the lack of a randomised study design. Four randomized trials (of which one is only available in Portuguese language and one only published in abstract form) have compared (both mechanical and balloon) dilation with intralesional steroid injections vs dilation alone. Two trials showed a significantly increased dilation free interval for the steroid arm and one showed an increase in stricture diameter in the steroid arm. In one study only 13% needed a re-intervention due to recurrent dysphagia in the steroid group whereas 60% needed re-intervention in the control group. This study included only patients with refractory peptic strictures, whereas the two other trials included patients of all sorts of strictures and patients with corrosive strictures, respectively. This was also the only study, in which a

standardized technique and symptom scoring system was applied and blinding of patients and effect investigators was adequately performed. However, a trial including patients with anastomotic strictures alone has not yet been performed.

Objective: To compare Savary dilation with saline 0.9% injections (placebo) with Savary dilation with triamcinolon injections (80 mg) in patients with benign anastomotic esophageal strictures.

Study design: A multicenter double-blind, placebo controlled, randomized trial.

Study population: All patients with dysphagia grade 2-4 due to benign anastomotic strictures after esophagectomy with gastric tube and cervical anastomosis.

Intervention: Patients in treatment arm A will be treated with Savary Galliard dilation after four saline 0.9% injections of 0.5 ml and patients in treatment group B receive Savary dilation after four intralesional triamcinolon 0.5 ml injections 40 mg/ml in all four quadrants.

Main study parameters/endpoints:

1. Median time to repeat dilation ;
2. Success rate after 6 months follow-up (number of patients dysphagia free within 3 dilation sessions).

Nature and extent of the burden and risks associated with participation, benefit and group relatedness: Patients will be asked to fill in a questionnaire on their symptoms before first treatment, and patients will be contacted by phone after 1 week (for complications only) after 2 weeks, 1 month, three months and six months for another questionnaire. Furthermore, patients will document their dysphagia score daily during the first month, thereafter; they will document their dysphagia score once weekly. No additional hospital visits or blood samples are required. The potential benefit expected in the study arm will be a longer dilation free interval and less dilation sessions to achieve relief of dysphagia.

Doel van het onderzoek

By injection of triamcinolone before dilation we hypothesise that the time to repeat dilation is increased and the total number of dilation sessions is decreased as well.

Onderzoeksopzet

Follow-up at 1, and 2 weeks and 1, 3, and 6 months.

Onderzoeksproduct en/of interventie

Injection of placebo (saline 0.9%) vs injection of Kenacort (80mg) followed by Savaray dilation up to 16 mm.

Contactpersonen

Publiek

P.O. Box 85500, 3508 GA Utrecht
M.M.C. Hirdes
Department of Gastroenterology and Hepatology
University Medical Center Utrecht
Huispostnummer F 02.618
Utrecht 3584 CX
The Netherlands
+31 88 755 1309/ #4596

Wetenschappelijk

P.O. Box 85500, 3508 GA Utrecht
M.M.C. Hirdes
Department of Gastroenterology and Hepatology
University Medical Center Utrecht
Huispostnummer F 02.618
Utrecht 3584 CX
The Netherlands
+31 88 755 1309/ #4596

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Patients with dysphagia grade 2-4 (Atkinson Dysphagia score) after esophagectomy with gastric tube reconstruction and cervical anastomosis;
2. Signed informed consent.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Previous dilation;
2. Dysphagia due to (suspicion of) malignant tumour recurrence;
3. Patients unfit for upper endoscopy;
4. Active anastomotic leakage or infection;
5. Recent vaccination with '®alive' vaccine;
6. During the acute phase of viral, bacterial or fungal infections;
7. Known gastric or duodenal ulcers;
8. Previous allergic reaction to one of the substances of Kenacort;
9. A known infection with tropical worms (such as Strongyloide) or parasites;
10. Outpatient visit within 6 months at the treating endoscopist.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blindering:	Dubbelblind
Controle:	Placebo

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-02-2010
Aantal proefpersonen:	60
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies

Datum: 09-03-2010

Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL2119
NTR-old	NTR2236
Ander register	METC UMC Utrecht / CCMO : 09/319 / NL29249.041.09 ;
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A