

Contribution of the Cultural Formulation Interview by practice nurses mental health in diagnosis and treatment of non-Western patients in general practice.

Gepubliceerd: 19-04-2013 Laatste bijgewerkt: 18-08-2022

Anxiety, depression and MUPS are common among first-generation non-Western migrants. The problem with the doctor also concerns motivating and treating these patients. The hypothesis is that attention to the social and cultural context of the patient...

Ethische beoordeling	Positief advies
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON26878

Bron

NTR

Verkorte titel

CI-POH

Aandoening

depression, anxiety, MUPS, depressie, angststoornissen, SOLK

Ondersteuning

Primaire sponsor: Primary performer: NIVEL (Netherlands Institute for Health Services Research)

Secondary performer: PHAROS (knowledge and advisory centre health for migrants, refugees and people with limited health literacy)

Overige ondersteuning: ZonMw (Netherlands Organisation for Health Research and Development), programme Winst door Verschil

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Compliance: Proportion patients not finishing their treatment prematurely within 6 months. Measured through reports from practice nurses and electronic patient record (EPR).

Toelichting onderzoek

Achtergrond van het onderzoek

In collaboration with care providers and representatives of the target group will the existing short form of the CI be developed into a version that can be used in the primary care. Next, the study will examine the impact of the use of the CI by practice nurses mental health on diagnosis, confidence in care, adherence, patient satisfaction, and indirectly health gains among Turkish, Moroccan, Surinamese and Antillean / Aruban immigrants.

A mixed methods study involves primary health care providers (GPs, nurses) to adapt the short version of the CI for use in primary care and tested in a pilot. Then a clustered effect controlled study will be conducted with the use of trained practice nurses applying the CI. A descriptive process evaluation supplemented by focus groups finally offer points for successful implementation in primary care. Quantitative data are collected with standardized questionnaires as well as information from electronic patient records. Qualitative information is obtained from the registration process, intake reports, EPR, interviews and focus groups with POH-GGZ, general practitioners and a focus group of patients in the intervention group.

Doel van het onderzoek

Anxiety, depression and MUPS are common among first-generation non-Western migrants. The problem with the doctor also concerns motivating and treating these patients. The hypothesis is that attention to the social and cultural context of the patient might lead to better doctor-patient relationship and more effective treatment. This could be accomplished by the Cultural Interview (CI) in primary care by the practice nurse mental health.

Onderzoeksopzet

Primary: Compliance - timepoint 6 months;

Secondary direct: Trust in health care provider (Health Alliance Questionnaire) - timepoint

month 0 and 6; no-shows (EPR) - timepoint month 6;

Secondary Indirect: Physical and mental health status (SF12) - timepoint month 0 and month 6, medical consumption [consultations, time for consultation, treatment, referral to 2nd care mental-physical-welfare] (EPR) - timepoint month 6 ; level stepped care (EPR) - timepoint month 6.

Onderzoeksproduct en/of interventie

Intervention:

Use of the Cultural Interview by practice nurse mental health spread over 2 consultations. Further care as usual.

Controle group:

Usual intake and care.

Contactpersonen

Publiek

PHAROS

PO Box 13318
Karen Hosper
Utrecht 3507 LH
The Netherlands
+31 (0)30 2349800

Wetenschappelijk

PHAROS

PO Box 13318
Karen Hosper
Utrecht 3507 LH
The Netherlands
+31 (0)30 2349800

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Non-western migrant patients from Turkish, Moroccan, Surinamese of Antillian origin;
2. 18-65years;
3. Referred by general practitioner to practice nurse mental health.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Native patients or migrant patients not from Turkish, Moroccan, Surinamese or Antillian origin.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Niet-gerandomiseerd
Blindering:	Dubbelblind
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-09-2013
Aantal proefpersonen:	210
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies

Datum: 19-04-2013
Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL3791
NTR-old	NTR3964
Ander register	ZonMw : 417100005
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A