# Touch ANd Go

Gepubliceerd: 04-08-2016 Laatst bijgewerkt: 18-08-2022

In human infants, tactile manoeuvres (warming, drying and rubbing the back or the soles of the feet) to stimulate breathing have been recommended during the initial assessment of the infant at birth (Lee 2011). These interventions alone would help...

| Ethische beoordeling | Positief advies       |
|----------------------|-----------------------|
| Status               | Werving gestopt       |
| Type aandoening      | -                     |
| Onderzoekstype       | Interventie onderzoek |

# Samenvatting

#### ID

NL-OMON27260

**Bron** Nationaal Trial Register

Verkorte titel TANGO

#### Aandoening

resuscitation spontaneous breathing preterm infant

## Ondersteuning

**Primaire sponsor:** Leiden University Medical Center (LUMC) **Overige ondersteuning:** Chiesi Pharmaceutical, LUF/Den Dulk Moerman Fonds

#### **Onderzoeksproduct en/of interventie**

#### **Uitkomstmaten**

#### Primaire uitkomstmaten

The main study parameter is the average respiratory minute volume at 1-4 minutes after birth (from 60 seconds until 240 seconds after birth).

# **Toelichting onderzoek**

#### Achtergrond van het onderzoek

Rationale Repetitive tactile stimulation can increase the respiratory drive/effort in preterm infants at birth, potentially leading to less CPAP failure and less positive pressure ventilation would be needed that might be injurious.

Objective To compare the direct effect of repetitive tactile stimulation on the respiratory effort of preterm infants during stabilisation at birth.

Study design A single blinded randomized clinical trial.

Study population Preterm infants of 27-31 weeks of gestation.

Intervention Infants will be randomized to receive repetitive stimulation during stabilisation after birth or standard stimulation. Repetitive tactile stimulation is hereby defined as gentle rubbing of the back and the soles of the feet for every 10 seconds as soon as the infant is placed on the resuscitation table. To prevent that the stimulatory effect will extinct, every 10 seconds of stimulation will be followed by 10 seconds without stimulation. Standard stimulation is defined according to the international guidelines, where stimulation is recommended initially if breathing is insufficient or absent.

Main study parameters/endpoints The main study parameter is the average respiratory minute volume at 1-4 minutes after birth (from 60 seconds until 240 seconds after birth).

Nature and extent of the burden and risks associated with participation, benefit and group relatedness The research is group-related, because most preterm infants need respiratory support at birth. The most logical way to avoid potentially harmful ventilation is to stimulate and support spontaneous breathing at birth. Tactile stimulation has been recommended during the initial assessment but infants could benefit from repetitive stimulation of the respiratory effort. Repetitive tactile stimulation is non-invasive and when applied gentle risks are negligible.

#### Doel van het onderzoek

In human infants, tactile manoeuvres (warming, drying and rubbing the back or the soles of the feet) to stimulate breathing have been recommended during the initial assessment of the infant at birth (Lee 2011). These interventions alone would help 10% of all infants that need

assessment after birth to achieve spontaneous breathing (expert's opinion) (Wall 2009). Although this is commonly accepted intervention, there are no human studies demonstrating the effect of stimulation on breathing at birth, especially in preterm infants. Because it is assumed that tactile stimulation during initial assessment promotes breathing, it is currently recommended in the international resuscitation guidelines (WHO 2012).

Currently, preterm infants are not dried, but placed in a plastic wrap to prevent hypothermia which may result in less stimulation (Rohana 2011, Morley 2007). Hereby, retrospective analysis of video recordings of stabilisation of preterm infants showed that tactile stimulation is often not performed. It is possible that applying stimulation repetitively augments the respiratory effort at birth and decreases the chance of CPAP-failure and the need for positive pressure ventilation.

#### Onderzoeksopzet

The first 10 minutes after birth.

#### **Onderzoeksproduct en/of interventie**

When an infant is included in the study, but the research protocol characterized for the allocated intervention is not strictly followed, the infant will be excluded for analysis and another infant will be included.

Infants will also be excluded if they are found to have a congenital abnormality or condition that might have an adverse effect on breathing or ventilation, including: congenital diaphragmatic hernia, trachea-oesophageal fistula or cyanotic heart disease.

## Contactpersonen

#### **Publiek**

LUMC Janneke Dekker Albinusdreef 2 Leiden 2333 ZA The Netherlands 071-5266641

## Wetenschappelijk

LUMC Janneke Dekker Albinusdreef 2 Leiden 2333 ZA

## **Deelname eisen**

## Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Preterm infants of 27-31+6 weeks of gestation can be randomized for receiving recurrent tactile stimulation or not.

## Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

When an infant is included in the study, but the research protocol characterized for the allocated intervention is not strictly followed, the infant will be excluded for analysis and another infant will be included.

Infants will also be excluded if they are found to have a congenital abnormality or condition that might have an adverse effect on breathing or ventilation, including: congenital diaphragmatic hernia, trachea-oesophageal fistula or cyanotic heart disease.

# Onderzoeksopzet

## Opzet

| Туре:            | Interventie onderzoek   |
|------------------|-------------------------|
| Onderzoeksmodel: | Parallel                |
| Toewijzing:      | Gerandomiseerd          |
| Blindering:      | Open / niet geblindeerd |
| Controle:        | Actieve controle groep  |

#### Deelname

| Nederland               |                 |
|-------------------------|-----------------|
| Status:                 | Werving gestopt |
| (Verwachte) startdatum: | 01-09-2016      |

4 - Touch ANd Go 31-05-2025

Aantal proefpersonen: Type:

Werkelijke startdatum

44

# **Ethische beoordeling**

Positief advies Datum: Soort:

04-08-2016 Eerste indiening

# Registraties

## **Opgevolgd door onderstaande (mogelijk meer actuele) registratie**

Geen registraties gevonden.

## Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

## In overige registers

| Register       | ID             |
|----------------|----------------|
| NTR-new        | NL5755         |
| NTR-old        | NTR6021        |
| Ander register | METC : P16.072 |

# Resultaten