Prevention of the return of nephrotic syndrome in children by adding levamisole to standard prednisone treatment.

Het voorkomen van terugval van het nefrotisch syndroom bij kinderen door het toevoegen van levamisol aan de standaardbehandelng met prednison.

Gepubliceerd: 06-02-2018 Laatst bijgewerkt: 18-08-2022

Adjuvant therapy of levamisole to prednisolone prevents relapses in children with a first episode of idiopathic nephrotic syndrome.

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON27331

Bron Nationaal Trial Register

Verkorte titel LEARNS

Aandoening

Idiopathic nephrotic syndrome / Idiopathisch nefrotisch syndroom Relapse / Recidief Childeren / Kinderen

Levamisole / Kinderen

Ondersteuning

Primaire sponsor: Academic Medical Center (AMC) Postbus 22600 1100 DD Amsterdam The Netherlands +31 (0)20 566 9111 Overige ondersteuning: Dutch Kidney Foundation

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The occurrence of relapse within 12 months after first presentation. Relapse is defined as the recurrence of proteinuria (3+ urine dipstick or >200 mg/mmol creatinine) on three consecutive days.

Toelichting onderzoek

Achtergrond van het onderzoek

Idiopathic nephrotic syndrome (INS) is a relatively rare disease, predominantly in children and adolescents, with an estimated incidence around 1.5 cases per 100,000 children/year in the Netherlands (approximately 60 newly diagnosed cases a year). After initial treatment with corticosteroids, the vast majority achieves remission. Unfortunately, relapse rates are high (80%), resulting in repeated and high doses of corticosteroids that have major physical and psychological side effects. Therefore, INS with its high risk of relapses might significantly impair the health-related quality of life (HRQoL) of affected children and may lead to substantial parental stress.

Previous randomized controlled studies (RCTs) showed promising results when levamisole, an antihelminthic drug, was added as adjuvant therapy to corticosteroids in children with frequently relapsing INS (FRNS) in reducing the occurrence of relapses. Therefore, we hypothesize that adding levamisole to corticosteroids as initial therapy in children with a first episode of INS will prevent relapses. This is substantiated by the fact that 1) levamisole is a immunomodulator that has the ability to skew Th2 immune response toward the Th1 response and 2) INS is characterized by a skewing of the immune response into Th2. As such

levamisole may prevent relapses of INS by restoring that balance between Th1 and Th2.

In comparison to other steroid sparing drugs, levamisole does not induce immunosuppression. It knows only little adverse effects of which neutropenia (<1500/mm3) is the most common and most serious and for which regular testing is required.

In addition, the underlying causes of INS and the prognostic factors to estimate the risk of relapse in INS patients are poorly understood. Also, little is known about the mechanism of action, the pharmacokinetics (PK), and pharmacodynamics (PD) of levamisole in children. Therefore, the RCT will be extended with 1) HRQoL questionnaires, 2) PK/PD analyses of prednisolone and levamisole (as well as the feasibility of measurement of levamisole concentration in saliva), 3) biobanking for future research, 4) study on the pathogenesis of INS, and 5) the mechanism of action of levamisole.

Our primary objective is to investigate the effect of additional levamisole in comparison with placebo from 4 weeks to 6 months after the start of the first episode of steroid sensitive INS in children (age 2 – 16 years) on the occurrence of relapses within 12 months. We hypothesize that adding levamisole to standard therapy with corticosteroids prevents relapses. To test our hypothesis, we will conduct an international (the Netherlands and Belgium), multicentre, double blind, randomized, placebo-controlled clinical trial. If remission is achieved, patients will be randomized in a 1:1 ratio to either levamisole (study group) or placebo (control group). Study medication will be used on alternate days for 24 weeks.

Doel van het onderzoek

Adjuvant therapy of levamisole to prednisolone prevents relapses in children with a first episode of idiopathic nephrotic syndrome.

Onderzoeksopzet

Primary endpoint: 12 months after first presentation of INS.

Secondary endpoints: 24 months after first presentation of INS. Secondary endpoints are analysed as time to first occurrence and number of occurrences during 24 months.

Onderzoeksproduct en/of interventie

- Prednisolone schedule according to French protocol: tapering schedule of 18 weeks (in comparison to 12 weeks of prednisolone according to Dutch protocol 'Werkboek Kindernefrologie').

- Additional 6-month (24 weeks) treatment of either 2.5 mg/kg levamisole or placebo on alternate days.

Contactpersonen

Publiek

Acedemic Medical Center (AMC)

F. Veltkamp Postbuss 22600

Amsterdam 1100 DD The Netherlands

Wetenschappelijk

Acedemic Medical Center (AMC)

F. Veltkamp Postbuss 22600

Amsterdam 1100 DD The Netherlands

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Inclusion criteria for Immunomics/Biobank:

- Age 2 to 16 years.

- First episode of idiopathic nephrotic syndrome, confirmed by: hypoalbuminaemia <25 g/L; Proteinuria >200 mg/mmol creatinine; Complement C3 within normal range.

- Written informed consent.

Inclusion criteria for RCT:

- Steroid sensitive nephrotic syndrome (remission achieved after 4 weeks of oral treatment with prednisolone).

- Weight >9 kg

- Ability to swallow (placebo) 5 mg tablet of study medication in children <6 years of age.
- Negative pregnancy test in girls who are of childbearing potential.
- Absence of contraindication for levamisole: neutropenia <1500/mm3.
- Ability to comply to study protocol.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Exclusion criteria for Immunomics/Biobank:

- Age <2 years or >16 years.

- Previous episodes of INS.

Exclusion criteria for RCT:

- Steroid resistant nephrotic syndrome (persistent proteinuria after 4 weeks of oral treatment with prednisolone).

- Previous or current malginancy, diabetes mellitus type 1, current liver disease, and/or convulsions.

- Hypersensitivity to levamisole or one of its substances (lactose).

Onderzoeksopzet

Opzet

Type: Onderzoeksmodel: Interventie onderzoek Parallel

Toewijzing:	Gerandomiseerd
Blindering:	Dubbelblind
Controle:	Placebo

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-03-2018
Aantal proefpersonen:	92
Туре:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	06-02-2018
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register
NTR-new
NTR-old
Ander register

ID NL6826 NTR7013 Dutch Kidney Foundation : CP16.03

Resultaten