

Does treatment with rosiglitazone result in improved pancreatic beta-cell function as compared to glimepiride in metformin treated diabetes type 2 patients?

Gepubliceerd: 07-02-2006 Laatste bijgewerkt: 18-08-2022

By inducing a shift of fat out of the visceral compartment - among which the pancreas - into the subcutaneous compartment rosiglitazone results in improved pancreatic beta-cell function in type 2 diabetes patients, as compared to a...

Ethische beoordeling	Positief advies
Status	Werving tijdelijk gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON27919

Bron

Nationaal Trial Register

Verkorte titel

N/A

Aandoening

Type 2 diabetes.

Type 2 diabetes is a heterogeneous disorder involving varying levels of insulin insensitivity and impaired islet beta-cell function.

Ondersteuning

Primaire sponsor: Academic Medical Center - Amsterdam

Overige ondersteuning: N/A

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

The peak insulin concentrations during the hyperglycaemic clamp protocol.

Toelichting onderzoek

Achtergrond van het onderzoek

Study title:

Does treatment with rosiglitazone result in improved pancreatic β -cell function as compared to glimepiride in metformin treated diabetes type 2 patients?

Introduction:

Thiazolidinediones, a new class of insulin sensitizing agents, have been shown to induce a shift of fat out of the visceral compartment – among which the pancreas – into the subcutaneous compartment. This could also result in a restoration or preservation of endogenous insulin secretion capacity, loss of which is one of the fundamental defects in Type 2 diabetes. A recent study could not confirm this hypothesis, but various shortcomings in the design of this previous study can be noted, most notably a treatment period that is likely to have been too short, and the fact that patients were not using metformin, the standard treatment for type 2 diabetes.

Aim of the study:

To investigate the effect of rosiglitazone treatment on β -cell function in type 2 diabetes patients as compared to a sulfonylurea derivative, while both groups continue metformin treatment.

Design:

Twenty-two patients will be randomized to metformin with glimepiride 4 mg a day or metformin with rosiglitazone 8 mg a day.

Patients:

Eligible patients are those with Type 2 diabetes using metformin. Exclusion criteria are established coronary heart disease and previous use of a thiazolidinedione.

Measurements:

Patients will undergo a 200 min hyperglycaemic (aiming at 10 mmol/l) clamp with administration of glucagon-like peptide-1 (GLP-1) starting at 120 min (bolus injection of 4.5 pmol/kg followed by a continuous infusion of 1.5 pmol/kg/min until the end of the clamp) and an arginine (5 g) bolus at 180 min to elicit a further β -cell response. Twenty-six weeks later, the assessments will be repeated, again on metformin, other study medication taken until the morning before this assessment.

Outcome measures:

Primary outcome measure will be the peak insulin concentrations during the hyperglycaemic clamp protocol.

Burden for the participants:

The risk for participants is judged to be minor. Participation mainly requires an investment of time and undergoing insertion of the sensors and blood sampling.

8-Aug-2007: trial has stopped because of stop cause problems with inclusion of patients.

Doel van het onderzoek

By inducing a shift of fat out of the visceral compartment - among which the pancreas - into the subcutaneous compartment rosiglitazone results in improved pancreatic beta-cell function in type 2 diabetes patients, as compared to a sulfonylurea derivative, while both groups continue metformin treatment.

Onderzoeksopzet

N/A

Onderzoeksproduct en/of interventie

Patients will be randomized to 26 weeks of treatment with metformin with glimepiride 4 mg a day or metformin with rosiglitazone 8 mg a day.

Before the start of the treatment patients will undergo a 200 min. hyperglycaemic (aiming at 15 mmol/l) clamp with administration of glucagon-like peptide-1 (GLP-1) starting at 120 min. and an arginine bolus at 180 min. to elicit a further beta-cell response.

Twenty-six weeks later, the assessments will be repeated, again on metformin, other study medication taken until the morning before this assessment.

Contactpersonen

Publiek

Academic Medical Center, Department of Internal Medicine F4-257,
P.O. box 22660
S.G.H.A. Swinnen
Meibergdreef 9
Amsterdam 1100 DD
The Netherlands
+31 (0)20 5667836

Wetenschappelijk

Academic Medical Center, Department of Internal Medicine F4-257,
P.O. box 22660
S.G.H.A. Swinnen
Meibergdreef 9
Amsterdam 1100 DD
The Netherlands
+31 (0)20 5667836

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Informed consent form signed;
2. Type 2 diabetes patients, according to WHO criteria;

3. Age 18-70 years;
4. Use of metformin, at least 500 mg a day;
5. HbA1c > 7.0% inclusive when on metformin alone, or > 6.5 % when on combination therapy of metformin and a sulfonylureumderivative.

Use of a sulfonylureumderivative is allowed, with a wash-out period of four weeks before the first assessments.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Established coronary heart disease;
2. Previous use of a thiazolidinedione.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	Actieve controle groep

Deelname

Nederland	
Status:	Werving tijdelijk gestopt
(Verwachte) startdatum:	01-09-2004
Aantal proefpersonen:	22
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies

Datum:

07-02-2006

Soort:

Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL549
NTR-old	NTR605
Ander register	: N/A
ISRCTN	ISRCTN52245496

Resultaten

Samenvatting resultaten

N/A