SteepRamp validation & tendon structure in DM2.

Gepubliceerd: 11-02-2010 Laatst bijgewerkt: 18-08-2022

Physical exercise has a prominent role in the treatment of type 2 diabetes mellitus (DM2). This research project aims to improve exercise intervention programs in DM2 by validating a novel easy applicable exercise test for determining physical...

Ethische beoordeling Status	Positief advies Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Observationeel onderzoek, zonder invasieve metingen

Samenvatting

ID

NL-OMON28345

Bron Nationaal Trial Register

Verkorte titel SteepRamp validation & tendon structure in DM2

Aandoening

Diabetes Mellitus type 2

Ondersteuning

Primaire sponsor: Erasmus MC, Universitair Medisch Centrum Rotterdam
Afdeling Revalidatie geneeskunde / Sportgeneeskunde
Intern adres: Kamer H. 026
postbus 2040
3000 CA Rotterdam
Nederland
Overige ondersteuning: The Netherlands Institute for Sport and Physical Activity (NISB) / Nederlands Instituut voor Sport en Beweging (NISB)

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Substudy A:

Validating the supramaximal SteepRamp exercise test to determine physical fitness in type 2 diabetes patients by correlating the result with a conventional VO2max exercise test and determining the test-retest reliability.

>

Substudy B:

Comparing Achilles tendinosis incidence in type 2 diabetes patients with the incidence in matched controls.

Toelichting onderzoek

Achtergrond van het onderzoek

N/A

Doel van het onderzoek

Physical exercise has a prominent role in the treatment of type 2 diabetes mellitus (DM2). This research project aims to improve exercise intervention programs in DM2 by validating a novel easy applicable exercise test for determining physical fitness (substudy A) and investigating if advanced glycation endproducts (AGE) in the skin can predict tendon structure abnormalities that predispose tendinopathy (substudy B).

Onderzoeksopzet

Substudy A:

1. Test day 1:

A. Intake interview and examination: A sports physician will screen the patient for cardiovascular disease and musculotendinous complaints and/or injuries using a questionnaire and short clinical examination;

B. Anthropometry: Besides weight and length, body fat percentage will be estimated form skin fold measurements using the Durnin & Womersly method. Abdominal circumference will be measured at the level of the belly button;

C. Spiro-ergometric test: Performed on a cyclo-ergometer. After a 4 min. warming-up period the resistance will be increase using a 2 watt/6 sec or a 1,2 watt/6 sec. protocol for men and

women respectively, resulting in approx. 10 min of exercise with resistance. The test will end with a 5 min. recovery phase. During exercise and recovery VO2, ECG, blood pressure, oxygen saturation and the rate of perceived exertion (Borg score) will be monitored;

D. Fasting venous blood sample & urine test

blood: Hemoglobin, HbA1c, plasma glucose, HDL-C, LDL-C, T-Chol, triglycerides and free fatty acids.

Blood plasma (3 ml) will be collected into EDTA containing tubes and will be centrifuged for 4 min. at 4°C. Aliquots of plasma will be frozen immediately in liquid nitrogen and stored at -80°C until analysis.

Urine: microalbuminuria (urine);

E. Activity monitoring: The patient will receive a validated multi-sensor accelerometer (Actigraph®) [36] and is requested to log daily activity to estimated kcal expenditure over 7 days;

2. Test day 2:

A. Sub-maximal and SteepRamp test: Within 7-14 days a sub-maximal exercise test will be performed using the following protocol: 2 min at 0,5 watt/kg, 2 min at 1,0 watt/kg and 2 min at 1,5 watt/kg. Ater 30 min a SteepRamp test will performed (3 min. warming-up phase without resistance, 25 watt/10 sec. test phase, test termination at <60 RPM). Maximal workload and the rate of perceived exertion (Borg score) will be documented;

3. Test day 3:

A. SteepRamp and sub-maximal test: Within 7-14 days the two tests will be repeated in reversed order;

4. Test day 4:

A. Sit-to-Stand test: A standardized Sit-to-Stand test [55] will be performed. The time to complete the test will be recorded. The test is validated as a quadriceps strength test;

B. Isokinetic quadriceps strength: Quadriceps strength will be objectified with a (isokinetic) Biodex® measurement (5 repetitions at 60 gr/sec).

Substudy B:

Only for the cases and matched controls included in substudy B, for the control subjects this is the only test day.

Test day 5:

1. Skin auto-fluorescence measurement: Forearm skin auto-fluorescence will be measured using the AGE-reader® [51]. The measurement is completely automated and is obtained by

3 - SteepRamp validation & tendon structure in DM2. 15-05-2025

placing the forearm on the device. The measurement takes approx. 1 min. to complete;

2. UTC measurement: The measurement is completely automated and is obtained by placing the lower leg in the UTC device [53, 54]. A 10 MHz linear-array transducer (Smartprobe 10L5, Terason 2000, Teratech, USA) is moved along and perpendicular to the tendon's long axis over a distance of 9.6 cm. Images are collected at regular distances of 0.2 mm. Four echo-types were assigned, based on the stability of intensity and distribution in contiguous transverse images, namely: I) highly stable; II) medium stable; III) highly variable and IV) constantly low intensity and variable distribution. The measurement takes approx. 30 min. to complete;

3. Venous blood sample (only for control subjects) HbA1c. Blood plasma (3 ml) will be collected into EDTA containing tubes and will be centrifuged for 4 min. at 4°C. Aliquots of plasma will be frozen immediately in liquid nitrogen and stored at -80°C until analysis.

Onderzoeksproduct en/of interventie

Physical exercise has a prominent role in the treatment of type 2 diabetes mellitus (DM2). This research project aims to improve exercise intervention programs in DM2 by validating a novel easy applicable exercise test for determining physical fitness (substudy A) and investigating if advanced glycation endproducts (AGE) in the skin can predict tendon structure abnormalities that predispose tendinopathy (substudy B).

Substudy A:

To improve the result and adherence of an exercise intervention in DM2 a individual tailored program is advised. Until now only a elaborative and expensive spiro-ergometry is available to optimize a exercise program for the individual patient. A short supra-maximal exercise test (SteepRamp-test, 25 Watt/10 sec) is used in post-chemotherapy patients and shows accurate correlation with VO2max measured with spiro-ergometry. The SteepRamp test has been proven safe in heart failure patients. Hence it can be suggested that the SteepRamp-test is accurate and effective in estimating physical fitness in DM2 patients. The study targets to validate the SteepRamp exercise test for estimating the VO2max in DM2 patients. The secondary objective is to correlate these data to results from a sub-maximal exercise test (0,5-1-1,5 watt/kg.2 min), an activity monitor and an isokinetic guadriceps strength test. The secondary goal aims to investigate the SteepRamp test as a measure to estimate daily activity pattern and muscle strength and to determine reference values for the DM2 population. To achieve the studies objective 135 DM2 patients (age 30-80 yr, BMI 27-40 kgm/2) will be included and undergo a spiro-ergometry, a SteepRamp and the sub-maximal exercise test on separate days. Furthermore the subjects will wear an activity monitor for 7 days and perform an isokinetic strength test.

Substudy B:

The prevalence of tendinomuscular overuse injuries is one of the main reasons of premature termination of the exercise program in DM2. Inactivity is considered an important cause of musculotendinous deconditioning and subsequent injuries. However, advanced glycation endproducts (AGE) of collagen in the skin is a known manifestation of DM2 and can easily be determined using a skin autofluorescence test. This research hypothesizes that the glycation of collagen will also occur in the musculotendinous tissue predisposing tendinopathy in type 2 diabetes. Skin autofluorescence will be correlated with an ultrasonic tissue characterization (UTC) technique. To test the hypothesis a group of 30 subjects (aged 35-60 yr) of substudy A will be requested to undergo a skin autofluorescence and UTC test. This group will be compared with a control group, matched for age en gender. When the hypothesis can be confirmed, exercise intervention programs for DM2 patients can be incorporated with a tendon strengthening program if necessary.

Contactpersonen

Publiek

P.O. box 2040 D. Visser Rotterdam 3000 CA The Netherlands +31 (0)10 7031679

Wetenschappelijk

P.O. box 2040 D. Visser Rotterdam 3000 CA The Netherlands +31 (0)10 7031679

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Substudy A:

- 1. Type 2 diabetes mellitus >2 yr;
- 2. Age: 30-80 years;
 - 5 SteepRamp validation & tendon structure in DM2. 15-05-2025

- 3. BMI: 27-40 kg/m2;
- 4. Formal permission to participate in the study by signing an informed consent form.

Substudy B:

- 30 cases from substudy A will be included in substudy B.
- 1. Type 2 diabetes mellitus >2 yr;
- 2. Age: 35-60 years, (15 male, 15 female);
- 3. Formal permission to participate in the study by signing an informed consent form.
- 30 controls will be matched for gender and age.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Substudy A and B:

1. Cardiovascular disease such as heart failure, cardiac ischemia or serious peripheral vascular disease (e.g. claudicatio intermittens);

2. Serious orthopedic of neurological conditions precluding an exercise test or muscle strength test.

Substudy B: 1. Use of fluoroquinolones.

Onderzoeksopzet

Opzet

Type:

Observationeel onderzoek, zonder invasieve metingen

Onderzoeksmodel:	Cross-over
Toewijzing:	Niet-gerandomiseerd
Blindering:	Open / niet geblindeerd
Controle:	Geneesmiddel

Deelname

Nederland Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-04-2010
Aantal proefpersonen:	165
Туре:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	11-02-2010
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL2092
NTR-old	NTR2209
Ander register	MEC Erasmus MC : 2010-066
ISRCTN	ISRCTN wordt niet meer aangevraagd.

7 - SteepRamp validation & tendon structure in DM2. 15-05-2025

Resultaten

Samenvatting resultaten

N/A