

Effect of a single dose of 14.8 MBq/kg (0.4 mCi/kg) 90Y-ibritumomab tiuxetan ("Zevalin") following first-line R-CVP therapy in patients with follicular lymphoma on conversion rate assessed by FDG-PET and on stem cell mobilisation.

A Phase II clinical trial.

Gepubliceerd: 14-07-2009 Laatste bijgewerkt: 18-08-2022

Zevalin consolidation after first line therapy with R-CVP in patients with follicular lymphoma grade 1-3a, stage 2-4 induces conversion of PET positive PR to CR in more than 40% of patients.

Ethische beoordeling	Positief advies
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON28762

Bron

NTR

Verkorte titel

N/A

Aandoening

follicular lymphoma/folliculair lymfoom
first line therapy/eerstelijnsbehandeling
90Y-ibritumomab tiuxetan

Ondersteuning

Primaire sponsor: Vumc, hematology department

Overige ondersteuning: sponsor

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

To determine the conversion rate of PR to CR (i.e. PET negativity) after a single dose of 90Y-ibritumomab tiuxetan (a dose of 14.8 MBq/kg or 0.4 mCi/kg, max 1184 MBq or 32mCi) in patients with grade 1-3a, stage II, III or IV follicular lymphoma with PET-positive partial remission on PET-CT scan following first line R-CVP therapy.

Toelichting onderzoek

Achtergrond van het onderzoek

To determine the conversion rate of PR to CR (i.e. PET negativity) after a single dose of 90Y-ibritumomab tiuxetan (a dose of 14.8 MBq/kg or 0.4 mCi/kg, max 1184 MBq or 32mCi) in patients with grade 1-3a, stage II, III or IV follicular lymphoma with PET-positive partial remission on PET-CT scan following first line R-CVP therapy. Also, to compare biological characteristics of bone marrow and stem cell harvest before and after a single dose of 90Y-ibritumomab tiuxetan.

Doel van het onderzoek

Zevalin consolidation after first line therapy with R-CVP in patients with follicular lymphoma grade 1-3a, stage 2-4 induces conversion of PET positive PR to CR in more than 40% of patients.

Onderzoeksopzet

Conversion rate assessed by PET CT scan 3 months (and if still positive) six months after 90Y-ibritumomab tiuxetan. Stem cell mobilisation before and three months after 90Y-ibritumomab tiuxetan.

Onderzoeksproduct en/of interventie

A single dose of 90Y-ibritumomab tiuxetan (a dose of 14.8 MBq/kg or 0.4 mCi/kg, max 1184

MBq or 32mCi) in patients with grade 1-3a, stage II, III or IV follicular lymphoma with PET-positive partial remission on PET-CT scan following first line R-CVP therapy.

Contactpersonen

Publiek

VU Medical Center

Department of Hematology
 PK 2 BR 018
P.O Box 7057

M.J. Wondergem
Amsterdam 1007 MB
The Netherlands
+31 (0)20 4442604

Wetenschappelijk

VU Medical Center

Department of Hematology
 PK 2 BR 018
P.O Box 7057

M.J. Wondergem
Amsterdam 1007 MB
The Netherlands
+31 (0)20 4442604

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Histologically confirmed (according to WHO classification) CD 20 positive follicular lymphoma, grade 1, 2 or 3a, stage II, III or IV (with indication for treatment following local guidelines) of all FLIPI scores;
2. Age 18 years or older, for stem cell mobilisation analysis age 18-61 years;
3. Having received 6-8 courses of 1st line R-CVP therapy;
4. WHO performance status 0 to 2;

5. Life expectancy of at least 6 months;
6. PET positive partial remission on PET-CT scan after 6-8 R-CVP;
7. Absolute neutrophil count (ANC) $1.5 \times 10^9/l$ or higher;
8. Platelet count of $150 \times 10^9/l$ or higher;
9. Hb > 6 mmol/l, (transfusion is allowed to achieve this);
10. Less than 25% bone marrow involvement after 6-8 R-CVP, measured by bone marrow biopsy;
11. Written informed consent obtained according to local guidelines.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Any other anticancer treatment for NHL except the first line R-CVP;
2. Prolonged cytopenia during first line induction chemotherapy requiring more than 2 weeks delay due to insufficient bone marrow reserve;
3. Prior external beam radiotherapy to $> 25\%$ of active bone marrow. (involved field or regional);
4. Prior myeloablative therapy;
5. Patients who have not recovered from the toxic effects of the first line chemotherapy;
6. Presence of gastric, central nervous system or testicular localisation at first diagnosis;
7. Any other malignancy or history of prior malignancy except non-melanoma skin cancer or stage 0 cervical carcinoma within the past 10 years;
8. Patients with pleural effusion or ascites;
9. Patients with abnormal liver function (bili > 1.5 ULN or ALAT > 2.5 ULN);
10. Active uncontrolled infection;
11. Known diagnosis of HIV infection;
12. Patients with abnormal renal function: serum creatinine > 2.5 ULN;

13. Known hypersensitivity to murine antibodies or proteins;
14. G-CSF or GM-CSF therapy within 2 weeks (or 4 weeks if pegylated) prior to administration of 90Y-ibritumomab tiuxetan;
15. Female patients who are pregnant or breast feeding, or adults of reproductive potential not employing an effective method of birth control during study treatment and for at least 12 months thereafter;
16. Concurrent severe and/or uncontrolled medical disease which could compromise study participation;
17. Patients who received any investigational drugs or underwent surgery less than 4 weeks before entry in this study or who have as yet not recovered from the side-effects of such treatment;
18. Patients with a history of psychiatric illness or condition which could interfere with their ability to understand the requirements of this study;
19. Patients unwilling or unable to comply with the protocol.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	N.v.t. / één studie arm
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	15-08-2009
Aantal proefpersonen:	31
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies

Datum: 14-07-2009

Soort: Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL1798
NTR-old	NTR1908
Ander register	METC VU MC : 09/173
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A