

Effects of omeprazole on the pharmacokinetics of irinotecan in cancer patients.

Gepubliceerd: 18-01-2008 Laatste bijgewerkt: 18-08-2022

To investigate the influence of omeprazole on the metabolism and plasma pharmacokinetics (PK) of irinotecan and its metabolites in cancer patients.

Ethische beoordeling	Positief advies
Status	Werving gestopt
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON28883

Bron

NTR

Verkorte titel

OMEPIRI

Aandoening

Cancer.

Ondersteuning

Primaire sponsor: Erasmus MC - Afdeling Interne Oncologie

Overige ondersteuning: fund=initiator=sponsor

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Plasma pharmacokinetics (PK) of irinotecan and its metabolites.

Toelichting onderzoek

Achtergrond van het onderzoek

This single center, open-label crossover study intends to investigate the possible pharmacokinetic interaction of omeprazole and irinotecan. The study will be performed at the Erasmus MC, location Daniel den Hoed Cancer Center. A total of 14 evaluable patients will be treated. Based upon recently conducted retrospective and prospective studies of the pharmacokinetics of irinotecan as a function of BSA (body surface area), which showed that BSA-based dosing does not reduce interpatient variability in irinotecan pharmacokinetics and drug-associated toxicities (Mathijssen, JCO, 2002 & de Jong, CCR, 2004), we will administer a 600 mg flat-fixed dose of irinotecan (3-weekly, 90-minutes i.v.) to all included patients. Patients will be deemed evaluable when treated with two courses of irinotecan; one course without and one course with concomitant use of omeprazole, and complete pharmacokinetic sampling and toxicity assessment has been performed according to this protocol. All patients will receive their first course of irinotecan without concomitant omeprazole followed by a second course of irinotecan with concomitant use of omeprazole (Losec MUPS) 40 mg QD, starting at day 8 of the first course until the third day of the second course. Both courses of irinotecan will be preceded by a midazolam hydroxylation test, performed on day 0 of both courses.

Doel van het onderzoek

To investigate the influence of omeprazole on the metabolism and plasma pharmacokinetics (PK) of irinotecan and its metabolites in cancer patients.

Onderzoeksopzet

N/A

Onderzoeksproduct en/of interventie

In this open-label crossover pharmacokinetic study, we will compare the plasma pharmacokinetics of irinotecan and its metabolites in patients treated with courses of irinotecan with and without concomitant use of omeprazole. Patients will receive irinotecan at 600 mg (90 min. i.v. infusion) without omeprazole during the first treatment cycle. In the second treatment cycle, these patients will be pretreated with 40 mg omeprazole for 14 days, and will then receive the second infusion of irinotecan (600 mg). Omeprazole will be continued until the third day after the second irinotecan infusion. Both courses of irinotecan will be preceded by a midazolam hydroxylation test, performed on day -1 of both courses.

Contactpersonen

Publiek

Daniel den Hoed Oncologisch Centrum, Afdeling Interne Oncologie, kamer AS-24. Groene Hilledijk 301

J.M. Bol, van der
Rotterdam 3075 EA
The Netherlands

Wetenschappelijk

Daniel den Hoed Oncologisch Centrum, Afdeling Interne Oncologie, kamer AS-24. Groene Hilledijk 301

J.M. Bol, van der
Rotterdam 3075 EA
The Netherlands

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Histological or cytological confirmed diagnosis of any form of (irresectable and/or metastatic) cancer, which is thought to be sensitive to irinotecan-treatment;
2. Age $\geq$ 18 years;
3. WHO performance status $\leq$ 1;
4. Adequate hematological functions (ANC $>$ $1.5 \times 10^9/L$ (ANC between 1.5 and 2.0, must be approved by the study coordinators), platelets $>$ $100 \times 10^12/L$);
5. Adequate renal and hepatic functions (serum creatinin $<$ 1.25xULN, bilirubin $<$ 1.25xULN; ALAT and ASAT $<$ 2.5xULN, in case of liver metastasis $<$ 5xULN; alkaline phosphatase $<$ 5xULN; gamma-GT $<$ 5xULN);
6. Written informed consent;
7. Complete initial work-up within two weeks prior to chemotherapy.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Pregnant or lactating patients; patients with reproductive potential must use a reliable method of contraception (excluding oral contraceptives), if required;
2. Serious illness or medical unstable condition requiring treatment, symptomatic CNS-metastases or history of psychiatric disorder that would prohibit the understanding and giving of informed consent;
3. Time between last antitumor treatment and first day of irinotecan therapy less than 4 weeks, provided that the patient has recovered from relevant toxic effects;
4. Radiotherapy within the last 4 weeks before the first course, if more than 20% of the bone marrow area is involved;
5. Major surgery within 4 weeks before the first course (to be evaluated by an MD);
6. Unresolved bowel obstruction or chronic colic disease;
7. Unwillingness to abstain from grapefruit(juice), star fruit (carambola), (herbal) dietary supplements, herbal tea, herbals and over-the-counter medication (except for paracetamol and ibuprofen) during the study period (starting two weeks before the first midazolam hydroxylation test);
8. Unwillingness to change medication, or no adequate alternatives available, in case of (chronic) use of CYP3A and/or P-glycoprotein inhibiting or inducing medication, dietary supplements, or other influencing compounds during the study period (starting two weeks before the first midazolam hydroxylation test);
9. Unwillingness to change medication in case of use of midazolam, temazepam and/or diazepam during the study period (starting two weeks before the first midazolam hydroxylation test);
10. Use of omeprazole or any other proton pump inhibitor during the study period (starting two weeks before the first midazolam hydroxylation test).

Onderzoeksopzet

Opzet

Type: Interventie onderzoek

Onderzoeksmodel:	Cross-over
Toewijzing:	Niet-gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving gestopt
(Verwachte) startdatum:	11-01-2008
Aantal proefpersonen:	14
Type:	Werkelijke startdatum

Ethische beoordeling

Positief advies	
Datum:	18-01-2008
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL1137
NTR-old	NTR1179
Ander register	METC : METC-2007-380
ISRCTN	ISRCTN wordt niet meer aangevraagd

Resultaten

Samenvatting resultaten

N/A