Eye Movement Desensitization and Reprocessing (EMDR) in children with a medically related trauma

Gepubliceerd: 03-05-2016 Laatst bijgewerkt: 15-05-2024

1) EMDR will lead to significant improvement of psychosocial functioning, quality of life and somatic complaints in children with elevated PTSD symptoms (subclinical/partial PTSD). 2) Children with unfavourable predictors (low socioeconomic status...

Ethische beoordeling	Positief advies
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON29358

Bron Nationaal Trial Register

Aandoening

EMDR, PTSS, PTSD, medical trauma, trauma, congenital heart disease, emergency department, aangeboren hartafwijking, SEH

Ondersteuning

Primaire sponsor: ErasmusMC Sophia Children's Hospital Rotterdam, the Netherlands **Overige ondersteuning:** InnovatieFonds Stichting Hartekind Vereniging EMDR Nederland

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Toelichting onderzoek

Achtergrond van het onderzoek

Around 1 in every 10 children admitted to the hospital for undergoing an invasive and painful medical procedure develops a posttraumatic stress disorder (PTSD) and circa 3 in every 10 children develop a partial/subclinical PTSD. Although PTSD is associated with unfavourable medical and mental health outcomes, children are often not structurally screened on PTSD. If PTSD symptoms are left untreated, this may harm the child's quality of life and psychosocial functioning. Currently, cognitive behavioural therapy (CBT) is the most evidence-based intervention for PTSD. A new innovative treatment is Eye Movement Desensitization and Reprocessing (EMDR), which is less intensive, is less burdensome for patients, works faster and is thus cheaper compared to CBT. This is the first RCT to investigate the direct effectiveness of standardized EMDR in children with medically related trauma. Therefore we will include all consecutive patients who underwent one or more admissions to the ErasmusMC Sophia Children's Hospital and who are 6-16 years old between July 2016 – July 2017, encompassing:

1) previously healthy children (with no underlying illness/handicap) admitted to the hospital for the first time after an unexpected emergency with no comorbidity (trauma type I) and

2) patients with congenital heart disease admitted for a recurrent procedure (trauma type II; e.g. surgery).

Patients with PTSD symptoms are allocated to: a) EMDR or b) care-as-usual CAU. All patients will receive adequate medical care (CAU).

Doel van het onderzoek

1) EMDR will lead to significant improvement of psychosocial functioning, quality of life and somatic complaints in children with elevated PTSD symptoms (subclinical/partial PTSD).

2) Children with unfavourable predictors (low socioeconomic status, less adequate coping) will benefit more from EMDR as compared to children with favourable predictors.

Onderzoeksopzet

Pre-post assessments (T1,T2) and a 6 months follow-up assessment (T3)

Onderzoeksproduct en/of interventie

2 - Eye Movement Desensitization and Reprocessing (EMDR) in children with a medicall ... 14-05-2025

Eye Movement Desensitization and Reprocessing (EMDR). EMDR is a standardized treatment method for trauma, consisting of a structured nine-phase approach to address the past, present, and future aspects of a traumatic or distressing memory. The EMDR treatment will consist of six weekly sessions of 1 hour.

In this study the active control group will receive care as usual, which means they get the regular medical care.

Contactpersonen

Publiek

Wytemaweg 8, kamer kp-2841

Maya Meentken Rotterdam 3015CN The Netherlands 010-70-32140

Wetenschappelijk

Wytemaweg 8, kamer kp-2841

Maya Meentken Rotterdam 3015CN The Netherlands 010-70-32140

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

The target group of this study consists of:

1) children and adolescents (6-16 years old) suffering from a partial (subclinical) PTSD due to one, sudden and unforeseen event, that occurred at least 4 weeks up to maximally three years ago, such as a traffic accident or severe acute illness. We include children who -after consultation at the emergency department at our hospital-, were hospitalized for the first

3 - Eye Movement Desensitization and Reprocessing (EMDR) in children with a medicall ... 14-05-2025

time and underwent invasive diagnostic procedures and/or treatment; and

2) children and adolescents (6-16 years old) suffering from (subclinical) PTSD due to invasive, repeated traumatic events of which these children/adolescent's know these events may happen again. We include children/adolescents with congenital heart disease, who underwent a repeated invasive medical procedure or examination (e.g. heart surgery or heart catheterization), at least 4 weeks up to maximally 3 years ago.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- 1) Mental retardation.
- 2) Inability to read or write Dutch.
- 3) Previous treatment for PTSD.
- 4) A score below the 60th percentile on the SVLK

5) A clinical score on the Clinician Administered PTSD Scale for Children and Adolescents (CAPS-CA)

Onderzoeksopzet

Opzet

Туре:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blindering:	Enkelblind
Controle:	Geneesmiddel

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-07-2016
Aantal proefpersonen:	128
Туре:	Verwachte startdatum

4 - Eye Movement Desensitization and Reprocessing (EMDR) in children with a medicall ... 14-05-2025

Ethische beoordeling

Positief advies Datum: Soort:

03-05-2016 Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 47373 Bron: ToetsingOnline Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL5665
NTR-old	NTR5801
ССМО	NL56156.078.16
OMON	NL-OMON47373

Resultaten