

Protection against HPV

Gepubliceerd: 04-08-2014 Laatste bijgewerkt: 15-05-2024

Antibody responses after a two-dose HPV vaccination schedule are non-inferior to a three-dose schedule. Level of two-dose schedule antibodies remains above plateau.

Ethische beoordeling	Positief advies
Status	Werving nog niet gestart
Type aandoening	-
Onderzoekstype	Observationeel onderzoek, zonder invasieve metingen

Samenvatting

ID

NL-OMON29611

Bron

NTR

Verkorte titel

HPV-2D

Aandoening

Vaccination, HPV, Immunogenicity, Two-dose schedule

Vaccinatie, HPV, immunogeniciteit, twee-doses schema

Ondersteuning

Primaire sponsor: National Institute for Public Health and The Environment, Bilthoven, the Netherlands

Overige ondersteuning: Dutch Ministry of Health, Welfare and Sports

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Cohort 1997-2000

- Antibody responses of vaccine-induced types after a two dose schedule up to 4 1/2 years

after the first dose.

Cohort 2001

- Level and kinetics of vaccine-induced antibody response after a two dose schedule at approximately 7,12 and 24 months.

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale:

The change from three- to two-dose schedule for HPV vaccination asks for monitoring of the kinetics of vaccine induced antibodies over time and of quality of vaccine-induced antibodies and cellular immunity after a two-dose schedule.

Objective:

- To study the level and quality of antibody response at approximately 7, 12 and 24 months following the first dose of HPV-16/18 vaccination in a two-dose schedule and check whether this level remains above plateau.
- To study whether antibody responses involved in a two-dose HPV-16/18-vaccination schedule compared to a three-dose schedule, are non-inferior at approximately 1 ½, 2 ½, 3 ½ and 4 ½ years after the first dose.

Study design:

In a prospective cohort study cellular immunity, the level and quality of vaccine-induced antibodies will be studied in girls born in 2001 who were vaccinated by a two-dose schedule in 2014. Cross-sectional observational sampling will be performed among girls born between 1997 and 2000, to compare the vaccine-induced antibody levels and avidity after a two-dose schedule with a three-dose schedule.

Study population:

Girls born in 2001 who received a two-dose schedule and girls born between 1997 and 2000

who received either two or three doses of the bivalent HPV vaccine.

Main study parameters/endpoints:

- Type specific antibody levels against HPV types 16,18 in serum following the two-dose schedule and whether these levels remains above plateau for HPV-16/-18 up to 24 months after the first dose
- Kinetics of type specific antibody levels against HPV types 16,18 in serum following the two-dose schedule up to 24 months after the first dose
- Whether the two-dose schedule is non-inferior with regard to HPV16/18 antibody levels to the three-dose schedule up to approximately 4 ½ years after the first dose

Doel van het onderzoek

Antibody responses after a two-dose HPV vaccination schedule are non-inferior to a three-dose schedule.

Level of two-dose schedule antibodies remains above plateau.

Onderzoeksopzet

Cohort 1997-2000

- 18, 30, 42, 54 months after the first dose

Cohort 2001

- 7, 12 and 24 months after the first dose

Onderzoeksproduct en/of interventie

None

Contactpersonen

Publiek

National Institute for Public Health and the Environment (RIVM)

Centre for Infectious Disease Control

Epidemiology and Surveillance Unit

PObox 1 (Pb.75)
Robine Donken
Bilthoven 3720 BA
The Netherlands
+31 (0)30 2744378

Wetenschappelijk

National Institute for Public Health and the Environment (RIVM)

Centre for Infectious Disease Control

Epidemiology and Surveillance Unit

PObox 1 (Pb.75)
Robine Donken
Bilthoven 3720 BA
The Netherlands
+31 (0)30 2744378

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

In order to be eligible to participate in this study, a subject must meet all of the following criteria:

- Vaccinated with the bivalent HPV vaccine (Cervarix)
- Received two- (with at least five months interval) or three-doses (0,1,6 months) of the vaccine
- Born between 1997 and 2001
- Female

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

A potential subject who meets any of the following criteria will be excluded from participation in this study:

- Unknown or incorrect address
- Deceased
- Participated in tolerability study and stated to be no longer approachable for further research

Onderzoeksopzet

Opzet

Type:	Observationeel onderzoek, zonder invasieve metingen
Onderzoeksmodel:	Parallel
Toewijzing:	Niet-gerandomiseerd
Blinding:	Enkelblind
Controle:	Geneesmiddel

Deelname

Nederland	
Status:	Werving nog niet gestart
(Verwachte) startdatum:	01-09-2014
Aantal proefpersonen:	418
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	04-08-2014
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 53042

Bron: ToetsingOnline

Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL4578
NTR-old	NTR4719
CCMO	NL48754.029.14
OMON	NL-OMON53042

Resultaten